November 16, 1948

Mr. U. E. Bowes
Owens-Illinois Glass Company
Toledo 1, Ohio

Dear Mr. Bowes:

Enclosed you will find three copies of a report on the results of animal experiments with Kaylo dust. As is our custom, we have summarized briefly material previously presented, including our interim report dated October 30, 1947, and have given detailed discussion only of subsequent developments. When all experiments have been completed, we expect to prepare a final report which will include details of each phase so that all data will be available in one place. However, the experimental study of the effects of inhaled Kaylo dust on normal uninfected animals is now finished and conclusions expressed on that subject are final rather than tentative.

In the report issued one year ago, which describes the findings in animals that inhaled Kaylo dust for periods up to 30 months, the following tentative conclusion was made:

> "In consequence of the experimental studies with guinea pigs to determine the biological activity of Kaylo, it may be tentatively concluded that Kaylo alone fails to produce significant pulmonary damage when inhaled into the lung."

During the 30 to 36 months period, however, definite indication of tissue reaction appeared in the lungs of animals inhaling Kaylo dust and therefore, I regret to say, our tentative conclusion quoted above must be altered. In all animals sacrificed after more than 30 months of exposure to Kaylo dust unmistakable evidence of asbestosis has developed, showing that [illegible] inhalation is capable of producing asbestosis and must be regarded [illegible] hazardous material. It should be noted that since neither silicosis nor the diffuse pulmonary fibrosis caused by inhaled diatomaceous earth was observed, the quartz and diatomaceous earth components of the dust apparently do not produce their typical lesions.

In order to present more information on the subject asbestosis, certain evidence derived from our experimental work with asbestos dust has been discussed. As these findings have not yet been released for publication, I request that, while using them as required in formulating a safety program, you regard them as confidential.

Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/1

Mr. [illegible]
Owens-Illinois Glass Company November 16, 1948

The new series of experiments with respect to the influence of Kaylo on tuberculous infection are well under way and are progressing satisfactorily. It is, of course, too early to expect significant results on which to base even tentative conclusions.

At this time may I review briefly the financial arrangement for conducting the investigation with Kaylo? The research program up to this year was carried on under a contract, initiated in 1945, by which the experiments would be subsidized with a grant of $5,000. per year. As pointed out in my letter of March 3, 1948, the contract terminated officially on February 15, 1948, but the investigation would be continued without charge until June because the original experiments were started late. Since a check was received which took care of the subsidy up to November 15, 1947, there is due on the old contract the sum of $1,250. for the final quarter of the contract (from November 15, 1947 to February 15, 1948). We have delayed sending an invoice for the final quarter until all work could be finished and a final report submitted. Following the termination of the experiment in June, it has required several months to do the histological work, study the tissue sections, collate the data and prepare the report which accompanies this letter.

We are including in the report a brief review of the new experiments in which the effect of inhaled Kaylo dust on tuberculous infection is being studied. Your purchase order AS-170 authorizing this experiment at $5,000. for one year is dated February 3, 1948, but owing to a shortage of animals and other unavoidable delays the actual experimental work did not get under way until May. Hence we have concluded that financial support for this new program should be dated from May 1, 1948. For reasons outlined in my letter of March 3, I suggested a two-year contract at $7,000. per year. Your letter of March 31 acknowledged [illegible] confirm the extra amount of $2,000. involved. Nevertheless, we have proceeded at the old rate and are endeavoring to absorb the increased cost from our Foundation reserve.

In a few days our accounting department will forward a statement of the payments now due in support of the research program, as follows:

Original experiment, begun February 15, 1945, at rate of $5,000. annually	
For final quarter (November 15, 1947 to February 15, 1948)	$1,250.
New experiment, begun May 1, 1948, at rate of $5,000. annually	
For first quarter (May 1, 1948 to August 1, 1948)	$1,250.
For second quarter (August 1, 1948 to November 1, 1948)	[illegible]
	$3,7[illegible]0.

Oct. 30, 1947 REPORT

Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/2

I realize that our findings regarding Kaylo are less favorable han anticipated. However, since Kaylo is capable of producing as- estosis, it is better to discover it now in animals rather than later n industrial workers. Thus the company, being forewarned, will be n a better position to institute adequate control measures for safe- uarding exposed employees and protecting its own interests.

Sincerely yours,

Arthur J. Vorwald, M.D.
Director

AV:LB
.cs. (3)

:: Mr. Dayton

Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/3

INTERIM REPORT

REGARDING THE BIOLOGICAL ACTIVITY

OF KAYLO DUST

to the

OWENS-ILLINOIS GLASS COMPANY

TOLEDO, OHIO

by

THE SARANAC LABORATORY

SARANAC LAKE, NEW YORK

OCTOBER 30, 1948

Submitted by:

[illegible]ald, M. D., Director
[illegible]dward L. Trudeau Foundation

In cooperation with:

Philip C. Pratt, M. D.
Thomas M. Durkan, M. D.
Anthony B. Delahant

cc: Aldrich
P.C. Pratt

Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/4

INTRODUCTION

One year ago, on October 30, 1947, an interim report was submitted, [illegible] of certain animal experiments with Kaylo over a total period of 30 months. It was necessary at that time to recommend further experiments concerning certain phases of the problem. The present report deals with the final phases of the experiment then in progress and with the early observations made in the new series of experiments.

The original experiment has now been completed and the conclusions drawn are based upon the results of a full three years' exposure to an atmospheric suspension of Kaylo dust. Previous experience has shown that this is the required period over which exposures to a dust of doubtful biological activity must be continued before final conclusions are warranted. This experiment, in fact, constitutes an example of the importance of such prolonged study for, as is to be shown below, the findings at three years were strikingly different from those at two and one-half years and the tentative conclusion reached at that time must now be altered. Thus, at the time of the previous report, the evidence indicated that Kaylo acted as an inert dust in normal rats and guinea pigs; resulting only in the formation of clumps of dust cells, without fibrosis. However, it will be shown below that Kaylo is capable, on prolonged inhalation, of producing asbestosis in the lungs of guinea pigs, and that it should be handled industrially as a hazardous dust.

This report will be presented in four sections as follows:

A. Final observations, dust inhalation in normal guinea pigs.

B. Discussion.

C. Conclusions.

D. Progress report on new series of experiments.

Arthur J. Vorwald to Owens-Illinois on Kaylo Causing Asbestosis in Animals (1948)
ACL Doc 11090 · Page 5 of 12
Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/5

Final observations, dust inhalation in normal guinea pigs.

Reference to the Interim Report of October 30, 1947, will reveal that two [illegible] pigs were placed in the dust room: one group in February and in August 1945. The animals were exposed to atmospheric suspensions of Kaylo dust for eight hours daily, five and one-half days a week throughout the experiment. The dust concentration, which varied somewhat from time to time, has averaged 116 million particles per cubic foot of air over the entire course of the experiment to date.

A brief review of the observations described on page 4 of the previous report will reveal that up to thirty months the reaction in guinea pigs was limited to the accumulation of clumps of macrophages, containing dust particles, in groups of adjacent air spaces. While occasional multinucleated giant cells were seen, there was no undue proliferation of tissue histiocytes and fibrosis was absent. A few unmistakable asbestosis bodies were identified. In general, the accumulation of cells of the types described above and their persistence as such over a full three year period, without the development of fibrosis, has been found to indicate that the agent responsible for them is not capable of producing fibrosis. Such agents are now considered to be "biologically inert," since the mere cellular accumulation has not been shown to affect pulmonary function or resistance to disease.

However, of nine animals sacrificed subsequent to thirty months in the present [illegible] have shown not only the above lesions but also have developed true fibrosis of a type characteristic of the response of guinea pigs to asbestos. This is a rather diffuse type of fibrosis which forms chiefly about the small air tubes, known as bronchioles, from which the air spaces, or alveoli, of the lung arise. The appended photomicrographs show the nature of the lesions being discussed. This condition is identical with asbestosis as can be seen by comparing

Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/6

ure 2 with figure 3.

The lesions in the tracheobronchial lymph nodes showed no change between [illegible] years.

[illegible] analysis of the lungs of guinea pigs that had inhaled Kaylo dust for periods up to 36 months yielded the data given in Table 1. It will be noted that as the period of exposure became longer the ash value gradually increased, indicating an accumulation of mineral matter in the lungs. There was a pronounced increase in the silica component up to about 30 months and then a slight decrease. The reason for this condition, which has been observed in experiments with other dusts, is not entirely clear. Since only two animals were analyzed for each exposure period, individual values were occasionally out of line but the general trend was clearly indicated.

Discussion.

The final observations in this experiment with Kaylo inhalation in normal guinea pigs, reported above and illustrated in the figures, prove that Kaylo is capable of producing the characteristic reaction, asbestosis. It is evident, therefore, that a considerable portion of the asbestos component in Kaylo remains unchanged during the manufacturing process or, if it is changed, that the altered product maintains its capacity to cause fibrosis. While the lesions up to 30 months showed no fibrosis, certain aspects of them were compatible with a preliminary [illegible] in the development of asbestosis. These aspects were masked by the short type of reaction to the materials other than asbestos in the dust. The characteristic early stage of the reaction to asbestos is the predilection of the fibers to localize in the lumen and wall of the small air ducts, known as terminal bronchioles. This is in contrast to particulate materials, which localize chiefly

the actual air spaces, or alveoli. In the present experiment the greatest [illegible] of the inhaled dust, being particulate, lodged in the alveoli and masked [illegible] action of some of the fibrous component. In retrospect, it is possible to [illegible] that there is more evidence of dust cell activity in the region of the bronchioles than is characteristic of the reaction to an inert particulate dust. Little significance was attached to this at the time of the previous report because there was no evidence of the development of fibrosis.

3. The differences between the early reaction to Kaylo and that to asbestos may be interpreted as a modification of the lesions by the particulate components of Kaylo. "Modification" of silicosis is well recognized when lesions of that disease develop following inhalation of a mixed dust containing quarts together with materials of an inert nature, such as iron oxide.

4. It is of importance to note that while the peribronchiolar reactions observed incriminate Kaylo as capable of producing asbestosis, they exclude the possibility that it may cause silicosis, since, over a full three year period, there is in no animal any focus of hyaline nodular fibrosis characteristic of the reaction to quarts. Furthermore, it may be added that the animals show no evidence of the type of diffuse fibrosis which has been observed in this laboratory to follow inhalation of diatomaceous earth dust. In regard to hydrated [illegible]-calcium silicate, a major constituent of Kaylo, it may be said that while there is no pointed evidence as to its effect on pulmonary tissue, there is much evidence to show that a variety of silicates, including some forms of calcium silicates, are biologically inert. It seems likely that the calcium silicate in Kaylo should conform with this previous experience.

5. Thus, the possibility expressed by Doctor Gardner in his early consideration of this problem, that Kaylo might produce two pneumoconioses, asbestosis and silicosis, with the resultant safety hazards, has been adequately studied, and concern over

Oct. 30, 1947

...er condition can be dismissed. Further consideration must be given to ... s, however.

...ently completed a large amount of work with asbestos, ... has been re... to certain other supporting companies, but not as yet ...blished. The following discussion of the problem is drawn largely from that ...rk, and we would request that these comments be considered confidential. It ...s felt that the following information may be of aid to the Owens-Illinois health ...partment in formulating a safety program, which certainly is necessary in view ...f the results of this Kaylo experiment.

Asbestosis, both in man and animals, is a chronic, slowly developing peri...nchiolar fibrosis, which in late stages extends from the original site of ...calization into the surrounding alveoli. The bronchiolar localization is appar...tly due to the fibrous nature of the mineral. Experimental studies indicate ...at the chief mode of action is by mechanical irritation of tissue by the fibers. ...e structure of these fibers, and the capability of repeated longitudinal divi...on to molecular strands, is important in the mechanism of action, since the ...lid fibers of glass wool are not capable of producing the fibrosis. Chemical ...ion is of little or no importance in causing reaction, since chrysotile rem...ved particulate by cutting and separation does not initiate fibrosis.

Certain investigations have indicated that a seemingly negligible proportion ...fib... asbestos is sufficient to produce the characteristic reaction. Thus, ...lation experiment was carried out with ball-milled asbestos in which an ...ttempt was made to eliminate all fibers from the material. It was found to be ...possible to break up all of the fibers and about one per cent of the air sus...pended dust consisted of fibers. The characteristic peribronchiolar fibrosis ...veloped in the exposed animals after forty months. Thus, it appears that very

Arthur J. Vorwald to Owens-Illinois on Kaylo Causing Asbestosis in Animals (1948)
ACL Doc 11090 · Page 9 of 12
Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/9

[illegible] numbers of fibers are capable of producing asbestosis, although the development of the lesions is delayed. The present experiment with Kaylo is also an [illegible].

[illegible] matter of progression of reaction after exposure to asbestos has been extensively investigated. The results indicate that in contrast to silicosis, where progression continues for some time, there is no significant degree of progression in extent of the lesions. Rather, the size of the lesions decreases in asbestosis due to the tendency of the fibrous tissue to contract after the stimulus to its production is removed. The irritation of tissue by an asbestos fiber is apparently arrested within a few months after it enters the lung by the formation of the "asbestosis body" (a coating of the fiber by iron-containing material), with the result that the existing fibrous tissue contracts. Whether this contraction exerts a beneficial or deleterious effect on lung function is now under investigation.

Unfortunately, the studies with asbestos have not as yet completely clarified the problem of its effect on associated tuberculous infection. It appears that Kaylo may exert a slightly more marked adverse influence on the infection than does pure asbestos. The difference may well be due to the presence of the small amount of quartz in Kaylo or perhaps to other incidental factors. One purpose of the studies now under way is to clarify this problem.

Conclusions.

1. Kaylo, because of its content of an appreciable amount of fibrous chrysotile, is capable of producing asbestosis and should be handled as a hazardous industrial dust.
2. Kaylo does not produce silicosis in animals.

[illegible]

Arthur J. Vorwald to Owens-Illinois on Kaylo Causing Asbestosis in Animals (1948)
ACL Doc 11090 · Page 10 of 12
Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/10

3. Kaylo does not produce the diffuse cellular fibrosis of the [illegible] observed as resulting from inhalation of diato- [illegible]

4. Progress Report on new series of experiments.

The three phases of the experiment, as outlined in the report of October 30, 1947 and in a letter dated March 3, 1948, are now under way. While little of significance has developed thus far, a review of the work is presented here. Progress and results to date of these experiments are given below under the title of each phase.

(1) The effect of Kaylo inhalation on tuberculous infection initiated at the onset of dust exposure.

Thirty experimental animals were infected with the attentuated strain (R_1) of tubercle bacillus used in this type of experiment and were placed in the dust room the next day. Thirty-three control animals were infected at the same time and kept in normal air. Of the experimental animals, six are now dead, two having been sacrificed at two months, two at four months and two having died.

One death occurred at five weeks, due to intercurrent pneumonia. The other was so recent that sections have not yet been prepared.

The animals killed at two months showed lesions comparable with [illegible] controls, both grossly and microscopically. The animals killed at four months have not been studied microscopically, but on gross examination the lesions appear similar in the two groups.

(2) The effect of Kaylo inhalation on regressing tuberculous lesions in guinea pigs.

In this phase of the experiment, twenty-five guinea pigs were infected, and groups of them were kept in normal air for two, four and six

Arthur J. Vorwald to Owens-Illinois on Kaylo Causing Asbestosis in Animals (1948)
ACL Doc 11090 · Page 11 of 12
Published by AsbestosClaims.Law
asbestosclaims.law/library/11090/11

[illegible] before being placed in the dust room. At the present time the first [illegible] have been placed in the dust room; the third is to go in on [illegible]

One, two animals of the first group have been sacrificed as yet, after two months in the dust. Microscopic observations have not been made, but in the gross, the lesions were comparable with those in control animals.

(3) The effect of accumulated Kaylo in the lungs on a subsequent tuberculous infection.

Thirty normal guinea pigs were placed in the dust room on May 12, 1948, and on August 12, 1948 were infected with the attenuated tubercle bacilli. Five animals were sacrificed at this time for chemical analysis to establish the amount of dust inhaled. Control animals were infected on the same date. Animals were sacrificed recently, two months after infection. Grossly, the lesions in the experimental and control animals were comparable, but again microscopic observations have not been made.

The results to date in this experiment are of little significance, since the disease is a chronic one and any changes due to the inhalation of dust of any nature are known to be slow. However, it can be said that the experiment is progressing satisfactorily, and that we anticipate none of the complications which interrupted the [illegible] station of the previous experiment in regard to the effect of Kaylo on experimental tuberculosis.