NHF

PLAINTIFF'S EXHIBIT

UC-3368

To (Name) Mr. J. F. Ballew, Jr.
Room Building 70-108A
Plant INSTITUTE PLANT

April 6, 1976

E.R. - Industrial Hygiene

Copy to Ms. B. J. Mitchell
Mr. J. G. Perkins
Mr. F. Williams

You have asked that measurements be made of personnel exposure to airborne asbestos fibers while weighing dishes are being prepared for residue analyses. Our records show that a similar measurement was made in Specifications Laboratory Room 205, March 10, 1975, and was reported to be slightly above threshold limit value (TLV).

Three determinations have recently been made by fiber count on filter samplers worn by employees while preparing sample dishes in the Analytical Laboratory Room 108. The results of these samplings are as follows:

Date	Employee	Fiber Count/ml Air
12/22/75	C. W. Devers	4.9
1/12/76	D. DeLuca	47.6
3/1/76	J. O. Good	7.7

We observed each of these samplings to be similar in technique and airborne dust was not visible in either instance. The TLV for asbestos dust exposure is currently 5 fibers greater than 5 micrometers in length per cc of air as a time weighted average (TWA) and a ceiling value not to exceed 10 fibers greater than 5 micrometers in length. As of July 1, 1976, the TLV will be reduced to 2 fibers as an OSHA allowable limit.

This job task is only infrequently performed and it is unlikely that the TLV as TWA would ever be exceeded. Our measurements, however, show that the ceiling level can be excessive. We, therefore, recommend that work involving preparation of sample dishes with asbestos fibers be performed within an exhaust ventilated hood.

H. H. Frazier

HHF:ls

404073

UCC 008858

Union Carbide trial exhibit UC-3368: H. H. Frazier to J. F. Ballew, Jr. on Airborne Asbestos Fiber Counts During Pre… Published by AsbestosClaims.Law
ACL Doc 18151 · Page 1 of 1 asbestosclaims.law/library/18151/1