Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle...
ACL Doc 18707 · Page 1 of 57

...ly one dose of serum prior to the onset ... received more than one dose as indicated in the ...ble. The dosage was usually 15 or 20 cubic centimeters.

Number of doses prior to onset of illness:	Number of patients
1	59
2	16
3	4
4	2
	81

The interval between the last dose of serum and the onset of illness ... as follows:

	Number of patients—		
	Receiving 1 dose	More than 1 dose	Total

AS A COURTESY WE HAVE PROVIDED THIS ARTICLE BECAUSE OF ITS LENGTH IN THE FORM OF TWO PAGES PER EXPOSURE, WITH A SPECIAL RATE. THIS DOES NOT OBTAIN EXCEPT IN SPECIAL CIRCUMSTANCES AND WILL BE SO TAGGED.

UCC 017675

A21509

U. S. TREASURY DEPARTMENT
PUBLIC HEALTH SERVICE

DEPOSITION EXHIBIT
22
PC-2
4-14-82

Public Health Bulletin No. 241
August 1938

A STUDY OF ASBESTOSIS IN THE ASBESTOS TEXTILE INDUSTRY

By

WALDEMAR C. DREESSEN, Passed Assistant Surgeon
J. M. DALLAVALLE, Passed Assistant Sanitary Engineer
THOMAS I. EDWARDS, Technical Editor
J. W. MILLER, Pathologist
R. R. SAYERS, Senior Surgeon

With the assistance of

H. F. EASOM, M. D., Director
M. F. TRICE, Engineer
Division of Industrial Hygiene
North Carolina State Board of Health

From the Division of Industrial Hygiene
National Institute of Health

PREPARED BY DIRECTION OF THE SURGEON GENERAL

PLAINTIFF'S EXHIBIT
UC-4095

UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON : 1938

For sale by the Superintendent of Documents, Washington, D. C. - - - - - - - - - Price 20 cents

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/1

ORGANIZATION OF THE NATIONAL INSTITUTE OF HEALTH

THOMAS PARRAN, *Surgeon General, United States Public Health Service*
L. R. THOMPSON, *Assistant Surgeon General, Director, National Institute of Health*

Division of Biologics Control.—Chief, Senior Surg. W. T. Harrison.
Division of Chemistry.—Chief, Prof. C. S. Hudson.
Division of Infectious Diseases.—Chief, Senior Surg. R. E. Dyer.
Division of Industrial Hygiene.—Chief, Senior Surg. R. R. Sayers.
Division of Pathology.—Chief, Surg. R. D. Lillie.
Division of Pharmacology.—Chief, Pharmacologist Director Carl Voegtlin.
Division of Public Health Methods.—Chief, Surg. J. W. Mountin.
Division of Zoology.—Acting Chief, Senior Zoologist, Willard H. Wright.
National Cancer Institute.—Chief, Pharmacologist Director Carl Voegtlin.
National Advisory Cancer Council.—Executive Director, Dr. Lewis Hinchey

(iii)

A2151C

UCC 017676

CONTENTS

	Page
Abstract	iii
Introduction	1
Objectives of the study	2
The working environment	3
Methods and instruments used in the study of the working environment	3
The manufacture of asbestos textiles	4
Preparation	5
Carding	6
Spinning	7
Winding, twisting, and spooling	8
Weaving	9
Brake-lined manufacture	10
British manufacturing methods	11
Occupational analysis and a description of the chief occupations	13
Preparation	15
Carding	16
Spinning	16
Spooling, twisting, winding, and treating	16
Cloth and tape weaving	16
All other	17
Results of the engineering study	17
Dust studies	17
Nature of dust found in asbestos textile factories	17
Composition	17
Particle size	18
Dust concentration in different occupations	18
Preparation	18
Carding	19
Spinning	19
Spooling, twisting, and winding	20
Broadcloth weaving	20
Tape, wick, and brake-lined weaving	20
Cord, rope, and braid weaving	21
Brake-lined finishing	21
Miscellaneous occupations	21
Previous dust exposures	22
Interpretation of dust counts	23
The control of asbestos dust	23
Preparation department	24
Willowing	24
Filing	25
Picking	25
Carding	26
Spooling and twisting	27
Weaving	28
Inspection	28
Summary of the study of the working environment	29

A21511

UCC 017677

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle… Published by AsbestosClaims.Law
ACL Doc 18707 · Page 3 of 57 **asbestosclaims.law/library/18707/3**

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle...
ACL Doc 18707 · Page 4 of 57
UCC 017678
Published by AsbestosClaims.Law
asbestosclaims.law/library/187074

	Page
Statistical description of employees	15
Occupational history	15
Comparison with other industrial workers	16
Medical study	48
Procedure	48
Past medical history	49
Symptoms of asbestosis	50
X-ray findings	54
Lung-field markings	54
Diaphragm	61
Physical findings	62
Chest expansion	63
Respiratory rate	65
Clubbing of fingers and curving of nails	66
Heart defects	66
Asbestos corns	68
Laboratory findings	69
Asbestosis bodies	69
Results of tests on urine	73
The diagnosis of asbestosis	74
Occupational history	75
X-ray findings	75
Classification of lung-field markings	75
Symptoms	77
Representative case histories	77
Incidence of asbestosis	89
Threshold concentration of asbestos dust	91
Incidence of asbestosis in different occupations	94
Asbestosis in Great Britain and the United States	95
Pathological findings in asbestosis	96
Gross appearance	96
Microscopic appearance	96
Selected autopsy reports	101
Acknowledgments	115
Summary of medical findings	116
Annotated bibliography	118

ILLUSTRATIONS

Figure	Page
1. Vacuum cleaner bag arrangement for collecting suspended dust samples	4
2. Flow sheet of American practice	5
3. Mill for crushing asbestos fiber	6
4. Fly opener or willower (note exhaust)	7
5. Exhausted carding machines	8
6. Spinning mules	9
7. Twisting machines	10
8. Exhausted braider	11
9. Rope making machine	12
10. Broadcloth weaving loom	12
11. Brake-band weaving loom showing three of the six sections with woven band	13
12. Flow sheet of British practice	14
13. Gross appearance of suspended dust collected with vacuum cleaner	21
14. Microscopic appearance of suspended dust collected with vacuum cleaner device at 200×	22
15. Exhausted piling bins	34
16. Exhaust ventilation used for piling operations (British practice)	35
17. Schematic drawing of exhaust system used with picker machine	36
18. Vertical exhaust ventilation applied to an unenclosed primary carding machine	36
19. Totally enclosed exhausted primary carding machine	37
20. Elevation and plan for spooling exhaust system	37
21. Exhausted spooling frame	38
22. Exhausted twisting frame	39
23. Exhausted broadcloth loom	40
24. Detail of exhausted loom shown in figure 23	41
25. Detail of exhaust used on British type loom	41
26. Exhausted brushing and inspection table	41
27. Form used for recording occupational history	45
28. Form used for recording past and present medical history	49
29. Percentage of persons, classified by dust exposure, who had certain disorders	52
30. Percentage of persons, classified by lung-field markings, who had certain disorders	53
31. Form used for recording X-ray findings	54
32. Scheme used in classifying X-ray markings	55
33. Percentage of males, classified by age, who had certain lung-field markings	56
34. Relation of ground-glass lung-field markings to length of employment	58
35. Relation of ground-glass lung-field markings to dust exposure	58
36. Incidence of ground-glass lung-field markings in relation to length of employment and dust concentration	59
37. Relation of diaphragmatic fixation to dust exposure. Three stages of diaphragmatic fixation are represented	60
38. Form used for recording physical findings	63
39. Photomicrographs of asbestosis bodies found in sputum. Enlarged 530 times except *B* which is enlarged 310 times. A scale ruled in units of 50 microns, has been drawn beside each asbestosis body	67
39a. Asbestos corn on thumb of 36-year old man employed 4 years in the card room of an asbestos textile factory	70
40. Portions of chest X-ray films which illustrate the successive changes brought about by long-continued inhalation of asbestos dust	76
41–51. Chest X-ray films illustrating individual case histories	78–88
52. Photomicrographs of lung sections showing the presence of nodular fibrosis and emphysema in a silicotic lung and diffuse fibrosis and emphysema in an asbestotic lung. A section of a normal lung is shown for comparison. Magnification 16×	98
53. Asbestosis body in intra-alveolar connective tissue. Practically no fibrosis in this area. Case H–21 Magnification 1,000×. (Army Medical Museum No. 52,255)	99
54. Asbestosis bodies accompanied by giant cells in an area of fibrosed lung tissue. Case H–21. Magnification 705×. (Army Medical Museum No. 52,255)	100
55. Case H–21. Right lung. Markedly thickened pleura with fusion of interlobar pleura. Dense fibrous tags over entire surface	102

Page

57. Case II-21. Left lung. Pleura much thickened. Fewer fibrous adhesions than were noted on the right lung..... 103
57. Case II-21. Section of right lung. Advanced asbestosis. Distended bronchi and air spaces at both apex and base. Diffuse fibrosis throughout lung substance and thickened pleura..... 101
58. Case II-21. Section of left lung. Distended bronchi and air spaces at apex. Rather uniformly distributed fibrous and emphysematous areas throughout rest of lung..... 105
59. Chest X-ray of case II-23, 15 months before death..... 107
60. Case II-23. Right lung. Entire pleural surface covered by dense fibrous tags..... 108
61. Case II-23. Left lung. Very few fine fibrous tags are noted on the pleura..... 109
62. Case II-23. Section of right lung. Moderately advanced asbestosis. Fibrous areas and emphysema uniformly distributed throughout the lung substance..... 110
63. Case II-23. Section of left lung. Shows less involvement than right lung..... 111
64. Case II-25. Chest X-rays taken 15 months and 1 month before death. Part of the difference between the two roentgenograms is due to a difference in X-ray technique..... 113
65. Case II-25. Section of right lung. Early asbestosis. Uniform distribution of fibrous areas throughout lung substance..... 114

UCC 017679

A21513

ABSTRACT

Asbestosis, a lung disease caused by long-continued inhalation of asbestos dust, was the principal physical defect found on medical examination of 541 men and women employed (or recently employed) in three asbestos textile factories. The primary effect of asbestos dust seems to be initiation of fine, interstitial, pulmonary fibrosis which was not observed to progress as far as the nodular stage. This fibrosis was observed in post-mortem examinations of three former asbestos workers. During life, lung fibrosis can be detected by X-ray examination. The more important symptoms of asbestosis are progressive dyspnoea, variable cough which sometimes raises blood-streaked sputum, and loss of weight. The processes that produce dust in asbestos textile factories were studied and recommendations for dust control are incorporated in this report. Enough dust counts (242) were made in all parts of these factories so that the dust exposure of each worker can be estimated. The percentage of persons in different occupational groups who were affected by asbestosis or any of its symptoms varied with the average dust concentration to which they were subjected and with their length of employment. The only cases of asbestosis, three in number, found below 5 million particles per cubic foot were diagnosed as doubtful; well-established cases occurred at higher concentrations. It appears from these data that if asbestos dust concentrations in the air breathed are kept below this limit new cases of asbestosis would not appear.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle...
ACL Doc 18707 · Page 5 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/5

A STUDY OF ASBESTOSIS IN THE ASBESTOS TEXTILE INDUSTRY

INTRODUCTION

Several important properties of asbestos have led to its widespread use in industry. As everyone knows, it is noncombustible and an excellent insulator against heat and electricity. Fewer people seem to be aware that its fibers are sufficiently long and flexible to be carded, spun, and woven like wool, flax, or cotton. Utilizing these properties of asbestos, industries have been developed in this country which employed more than 6,000 men and women in slightly more than 60 establishments in 1933.

Another property of asbestos has been discovered more recently. When asbestos dust is inhaled daily, year in and year out, a lung disease termed asbestosis may develop. In many respects asbestosis resembles silicosis, but it differs from silicosis in several important particulars which will be discussed in the medical section of this report.

The first record of a case of asbestosis seems to have been made by Montague Murray in 1900. The first complete description of the disease and of the "curious bodies" seen in lung tissue and sputum appeared in 1927 when Cooke (9) and McDonald (22) reported two cases of asbestosis and listed their reasons for believing that asbestos bodies originate from asbestos fibers that reach the lungs. The publication of their papers aroused general interest in the subject and numerous papers appeared soon afterward. Hoffman (16) appears to have been the first American to call attention to the magnitude of the asbestosis problem. In 1918 he reported that 13 deaths from asbestosis had occurred among asbestos textile workers, and about the same time Pancoast, Miller, and Landis (29) reported on 17 cases of asbestosis. Mill's (26) paper was the first pathological report on asbestosis published in the United States, and in the same year, 1930, Lynch and Smith (38) reported on asbestosis bodies found in the sputum of asbestos workers.

Because several satisfactory literature reviews of asbestosis have appeared in recent years there seems to be no need to present a complete literature review in this report. Some of the more useful literature reviews have been listed in the annotated bibliography. In Merewether's (25) review emphasis is laid on the relation of asbestosis

A21514

UCC 017680

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle… Published by AsbestosClaims.Law
ACL Doc 18707 · Page 6 of 57 asbestosclaims.law/library/18707/6

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle...
ACL Doc 18707 · Page 7 of 57

to working conditions. The clinical aspects of the subject are well treated. Gloyne's (16) discussion of the pathology of asbestosis is particularly helpful. Middleton's (17) Milroy lectures on pneumoconiosis contain a section on asbestosis. Egbert (13) has summarized the literature reports on 28 fatal cases of asbestosis. "Active pulmonary tuberculosis was present in 6 instances of the total 28 cases; the fatal outcome was primarily the result of tuberculosis in 3 instances."

OBJECTIVES OF THE STUDY

The Public Health Service was requested by the State Board of Health and the Industrial Commission (administrator of the Workmen's Compensation Act) of North Carolina to assist them in making an engineering and medical study of health hazards in the asbestos textile industry.[1] The objectives of this study were threefold: (1) To make a medical study of the effects of long-continued inhalation of asbestos dust on the human body; (2) to identify the manufacturing processes that create dust and to recommend practices and, when necessary, equipment that will reduce the dust exposure of workers; and (3) to find out what concentration of asbestos dust can be tolerated without injury to health.

[1] The project was one of the many uses of Social Security funds for a cooperative project carried on by the Public Health Service and a state health department.

UCC 017681

A21515

THE WORKING ENVIRONMENT

METHODS AND INSTRUMENTS USED IN THE STUDY OF THE WORKING ENVIRONMENT

The evaluation of the working environment and its relation to the health of workers forms the basis of all field investigations conducted by the Division of Industrial Hygiene. The procedure followed by this Division is described in Public Health Bulletin No. 217 (2). Therefore, only a brief discussion of the subject is presented in this report.

The study of an occupational dust problem begins with a preliminary survey designed to obtain data on all the occupations in a given plant. It includes the number of workers employed in each occupation and the materials used in each process. The survey also reveals the number of workers exposed to a given hazard. Following the preliminary survey an evaluation of the occupational environment is made. In the dusty trades this includes a determination of the amount of dust present in the air, its composition and particle size, and the measurement of the effectiveness of exhaust systems where they are used to control dust hazards. The latter, in this instance, involved airflow measurements of several exhaust systems by means of a pitot static tube. The effectiveness of these systems was determined by dust counts.

THE MANUFACTURE OF ASBESTOS TEXTILES

Approximately 90 percent of the asbestos used in these plants is obtained from Canada. The remaining 10 percent comes from Arizona or South Africa, and, infrequently, from Russian and Australian sources.

Much of the machinery that is employed for the manufacture of asbestos textiles is identical with the machinery used for the production of cotton and woolen goods. Crude asbestos fiber is sent first to the preparation department and then, in the order named, to the carding machines, spinning frames, winding bobbins, twisting machines, spooling frames, and finally to the looms and miscellaneous fabricating devices.

The four plants investigated in the present study fall naturally into two classifications. Two plants convert crude asbestos and raw cotton into asbestos yarn and woven goods, but do not prepare the

Published by AsbestosClaims.Law
asbestosclaims.law/library/187077

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle...
ACL Doc 18707 · Page 8 of 57

products for the ultimate consumer. The other two plants purchase both asbestos yarn and unfinished woven asbestos tape. These establishments weave brake bands and convert their own woven product, as well as purchased material, into finished goods. The practices followed in the plants studied are schematically presented in the flow sheet shown in figure 2.



A 21516
UCC 017682

FIGURE 1.—VACUUM CLEANER BAG ARRANGEMENT FOR COLLECTING SUSPENDED DUST SAMPLES.

PREPARATION


Crushing.—The first step in the manufacture of asbestos textiles is the crushing of the crude stock in a milling machine. This is accomplished by large iron rollers which revolve in a steel pan in which the crude material is placed. (See fig. 3.) This operation separates the fibers and loosens the impurities that survived the cobbing of the ore at the mines. The crushing operation does not produce much dust, the fibers being disintegrated by a mashing and twisting action.

Most crude ore must be crushed for 20 minutes, although in some cases it may be satisfactorily disintegrated in 5 minutes. The crushed asbestos has a matted appearance. Following the crushing, the asbestos is replaced by hand in the bags for delivery to the willower.

Willowing.—The willowing or opening of the asbestos is essentially a cleaning operation. The willower beats and shreds the matted

FLOW SHEET

AMERICAN PRACTICE



Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/8

asbestos masses into individual fibers and removes some of the rock. The asbestos fibers that are retained are delivered directly to a storage bin or conveyed to one by pneumatic duct. (See fig. 4.) The material falling to the floor is screened in order to separate the grit from the short fibers. The grit is sold to cement manufacturers and the short asbestos fibers to paint, paper, and roofing manufacturers.

The willower is fed manually by a laborer who lifts the crushed asbestos into the willower hopper by means of a pitchfork. The beating and combing action which takes place within the machine produces a large amount of dust which escapes from the loose-fitting enclosure. The production of dust is increased when waste asbestos, recovered from the cyclone separators which form a part of the dust-




FIGURE 3.—MILL FOR CRUSHING ASBESTOS FIBER.

removal system, is willowed for reclaiming purposes. This latter operation is called fly willowing.

At one of the plants studied, the willowed asbestos is passed over a vibrating screen. This operation insures a cleaner product, since additional grit and short fibers are removed. Where a second screen is not available the asbestos may be passed through the willower a second time, a practice which is followed at two of the plants studied. The screening of asbestos is a decidedly dusty operation.

Mixing.—The asbestos fibers consist of blends of various grades and often include a certain amount of waste recovered by the fly willower. This process is followed in order to obtain end products with the characteristics demanded by the market. When a batch of asbestos has been made, it is then mixed with cotton. Asbestos fibers must be mixed with cotton before they can be spun and woven. The amount of cotton used depends upon the quality of the willowed asbestos fibers used. As a rule, from 5 to 15 percent by weight of all the asbestos textiles produced in this country is cotton.

Piling.—At the plants studied, the mixing is accomplished in two steps, the first consisting of placing the cotton and asbestos in alternate layers in a pile upon the floor, and the second step of feeding the mixture into a picker machine. The batch making, or the placing of the cotton and asbestos, is known as piling. It is productive of much dust since it is necessary to spread out both the cotton and asbestos into uniform layers, which involves the use of a rake and much trampling of the materials.

Picking.—After the mixture is made, it is lifted into the breast-high hopper of the picker machine. The material emptied into the




FIGURE 4.—FLY OPENER OR WILLOWER (NOTE EXHAUST).

hopper displaces air in the hopper which causes dust and fibers to escape upward into the face of the worker. In addition, the cylinder and accessory rollers of the picker machine operate at a high speed and much fine material escapes from the many small openings in the frame. The dust that is fanned out of the machines, together with that produced in the feeding operation, makes the pickerman's occupation comparatively dusty. In some plants two pickers are operated in series and in others only one, through which the material may be passed several times. In either case the picker discharges directly or by pneumatic duct into a collector bin. Where only one machine is available, however, the pickerman must move the material by pitchfork from the collector bin to the floor of the picker room for its recirculation through the picker. At one of the plants studied there are two pickers operated in series and connected by pneumatic

UCC 017683
A 21617

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavalle...
ACL Doc 18707 · Page 9 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/9

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 10 of 57

ducts. Here the collector bin is also the storage bin. In order to prevent the mixed material from accumulating in a steep pile a man is stationed in the stock bin at all times. This occupation creates considerable dust, since the cotton-asbestos mixture is delivered into the bin by overhead duct discharge. In order to reduce the dust exposure of the stock-bin man, his position is made interchangeable with the pickerman, each of whom spends a half of every day in the stock bin and a half day at the picker. In the plant having only one picker, the mixture is trucked from the collector bin to the storage bin. At one of the plants studied, exhaust ventilation is provided for the picker machines and the mixed stock storage bins. Oil sprays are




FIGURE 5. EXHAUSTED CARDING MACHINES.

frequently used within the picking machines in an attempt to prevent the formation of dust.

CARDING

From the storage bins the stock, as it is called, is hand-trucked to the carding machines where it may be piled upon the floor or fed into the hopper of the breaker card.

The cards perform several duties. They give the fibers an additional mixing, arrange them in parallel positions, and form them into roving. The term roving is applied to the strands of parallel fibers that are produced. A carding unit consists of two almost identical machines. The one through which the cotton-asbestos mixture passes first is termed the breaker card, and the other, the finishing card. The two machines are quite similar in that a large revolving cylinder with several accessory rollers performs the major portion of the work. The carding operation is quite similar to that of the

UCC 017684

A21518

picker, the essential difference being that the roller and cylinder surfaces are studded with fine steel bristles rather than large hooks. Some of the rollers straighten the fibers by combing out the tangles, while others doff, or remove, them from the combing rollers and the cylinder. Aside from these main features, the two units differ slightly. The breaker card is equipped with a feeding hopper similar to that used on the picking machine, while the finishing card is provided with a device for forming the carded fibers into strands. In the plants studied a system of conveyor lattices transfer the carded fibers from one machine to the other. The outside strands of roving are returned




FIGURE 6.—SPINNING MULES.

to the picker for reworking. The remainder of the roving is wound on long rollers with flanged ends, called jack spools.

The bristle-studded cylinders of the carding machines revolve at high speeds and much lint is thrown into the atmosphere. The effect produced by the revolving cylinders and wheels is similar to that of a fan, and as a result much of the dust remaining on the stock at this point, as well as asbestos and cotton fibers, is blown into the air. All carding machines studied were provided with exhaust ventilation.

While practically all of the roving is spun into yarn, some of it is marketed without further processing other than winding on spools or into coils. In this form it is used for electrical insulation and for conversion into packing for machine bearings.

SPINNING

The jack spools containing the roving may be conveyed either to the mule spinning machines shown in figure 6 or to the ring spinning frames. The spinning, on whatever machine performed, is a twisting operation which imparts strength to the strand of fibers by inter-

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/10

locking or meshing them together. The ring spinner twists the roving as it is wound on a spindle, while the mule spinner performs the same treatment in two operations; first, spinning it in 5- or 6-foot lengths and then winding it on a spindle. The ring spinner makes a yarn that is stronger and better in many respects than that produced on the mule spinner.

WINDING, TWISTING, AND SPOOLING

The spun yarn is converted into stronger thread in two operations. It is taken first to the winding machines where it is transferred from spindles to tubes. After being wound on tubes, the yarn goes to the twisting machines where the thread from two or more tubes is brought together and twisted into one thread. In the twisting operation the thread is frequently strengthened with brass, zinc, or copper wire, or




FIGURE 7.—TWISTING MACHINE.

with cotton twine. Following the initial twisting operation the yarn may return to the winders for placing on special tubes, in which form it is marketed. Then again, it may be conveyed to the cop or twill winders where it is wound into spindles for use in the shuttles of the looms. There are a variety of winding operations that prepare the yarn in some particular manner for the market or for use on certain types of machines. Likewise, there are a variety of twisting and spooling operations. (See fig. 7.)

The winding and twisting of the yarn is followed by a number of more or less related operations. The thread may be converted into cloth, tape, wick, brake lining, tubes, braids, or formed into rope. (See figs. 8 and 9.) A small amount of the product of the winders and twisters is treated by passing through a sizing compound, and subsequently dried on reels before being wound again on tubes for the

WEAVING

The weaving of asbestos is done on looms that operate in the conventional manner. Both cloth looms and tape looms are used. (See figs. 10 and 11.) On the latter, brake lining, asbestos tape, and wick for oil stoves are woven. The tape looms differ from the cloth looms in that the warp is segregated into several strips of material and in that there is a shuttle for each strip of warp. The weaving of all strips proceeds simultaneously; one mechanism operates all shuttles and a single reed suffices for all warps. Looms for tape weaving are made in multiples of 6 spaces up to as many as 30. Both dry and wet weaving are employed for the production of both cloth and tape. In wet weaving the warp is passed through a shallow pan of water and the cop (for use in the shuttle) is dipped into water.




FIGURE 8.—EXHAUSTED BRAIDER.

Dry weaving of broadcloth produces much lint and dust. The threads of the warp are continually rubbed past each other as they are moved from positions above or below the line of the shuttle which is shot from one side of the loom to the other at each change in position of the warp. Fiber fragments are rubbed from the threads at each change in their position and the fanning of the air by the moving machinery carries them into the atmosphere in and about the loom. The same machines are used for both wet and dry weaving.

In the manufacture of asbestos woven products at some of the plants studied, the warp is formed on individual looms by harnessing the thread from many spools which are hung on creel racks. At others

UCC 017685

A21519

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 11 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/11

the warp is first wound on a beam from the individual spools and then transferred to the loom. The latter practice makes a more compact weaving unit as creel racks occupy several times as much floor space as the loom. The creel racks are attended by a worker who is kept




FIGURE 9.—ROPE-MAKING MACHINE.




FIGURE 10.—BROADCLOTH WEAVING LOOM.

busy tying broken threads and replacing empty spools with full ones. The occupation involves an exposure to asbestos dust for which the looms are largely responsible.

The asbestos cloth is usually shipped to firms that convert it into [illegible] and other products that require a non-

UCC 017686

A21520

combustible fabric. Some asbestos wick is packed for the market without further treatment. Asbestos tape is usually sold to other industries. Some plants that weave brake bands sell it as a crude woven ribbon of fabric to finishing factories, while other manufacturers of the material convert it into finished products.




FIGURE 11.—BRAKE-BAND WEAVING LOOM SHOWING TABLE OF THE SIX SECTIONS WITH WOVEN BANDS.

BRAKE-BAND MANUFACTURE

The manufacture of brake bands proceeds in several steps, the first being the impregnation of the woven fabric with either an asphaltic compound or a vegetable resin. The latter is the newer form of treatment and is rapidly supplanting the former. Following this operation a heat treatment is applied in a specially constructed oven. In practice the ribbon of woven asbestos passes first into a vat of the

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 12 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/12

treating substances and thence immediately into a baking oven. These operations are carried on in a building that is well ventilated and which is removed from other structures sufficiently to reduce the fire hazard. The treated asbestos fabric is now calendered into uniform width and thickness and one surface is ground smooth. The brake-band material is next cut into suitable lengths and holes drilled into them according to pattern. Finally, the finished brake bands are branded and packed for the market.

In some plants molded brake bands are made of asbestos yarn and




FIGURE 12.—FLOW SHEET OF BRITISH PRACTICE.

a synthetic resin. The process consists of passing asbestos thread through a bath of resinous material and immediately winding it upon a spool or cylinder. Enough of the resin will adhere to the asbestos to form a matrix into which the asbestos thread will be imbedded. When the desired thickness is obtained, the spool or cylinder is given a heat treatment which hardens the resin. The baked cylinder of material is removed from the spool and cut into narrow sections. The circular bands thus formed are ground, machined, and otherwise made

UCC 017687

A 21521

The grinding or surfacing of the brake bands produces considerable dust. In the plants studied exhaust ventilation was used to control the exposure.

BRITISH MANUFACTURING METHODS

British practice differs in many respects from American practice. The same type of machinery is used in both countries but in British factories, dust-control equipment seems to be more generally used than in American factories. British factories specialize in the production of fine yarn and this involves numerous differences in practice.

In England the processing of the asbestos begins, as it does in the United States, with the crushing of the ore, but when this operation has been completed the manufacturing process does not parallel American practice until the fiber has been carded and is ready for spinning into yarn. (See fig. 12.)

After crushing, various grades of asbestos are blended in a machine quite similar to the conventional type of rotary concrete mixer. It is of passing interest to note that the same device is used for blending tea. The blend of asbestos is passed through a magnetic separator where certain mineral impurities are removed. It is cleaned further on rotary shaker screens where some grit and fiber fragments are eliminated. After screening, the material is fed into a willower. The British employ a vertical willower that is known as a Creighton opener. Its action is similar to that of the American willower, and the internal revolving mechanism is retained. The willowed material is subjected to a final cleaning on a second set of screens; such an operation is termed "sieving." Willowing is the last of the cleaning processes, after which the asbestos is placed into bins and is thereafter known as stock. It is now fully prepared for carding.

The practice at the plants included in this investigation is to add cotton to the asbestos and then subject the combination to a thorough mixing in a picker machine. In England cotton is added to the asbestos on the carding machine, both fibers being fed separately into the breaker card. The cotton in all probability is prepared for the carding operation on a conventional type of combination cotton textile picker machine which opens the fibers, cleans them, and forms the material into laps. The latter term is applied to the cotton which is formed into a broad ribbon of felt and then wound into a roll.

English carding machines are equipped with a feeding conveyor, or lattice, upon which first the asbestos and then the cotton are placed. The asbestos is added from a weighing box that is supported over and near the outside end of the conveyor. This box feeds asbestos in accordance with a predetermined ratio of the fiber to the cotton. When a certain weight of asbestos has accumulated in the box, a mechanism opens the bottom and allows the material to fall upon the

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 13 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/13

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 14 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/14
UCC 017688
A 21522

lattice. Further along a roll of cotton felt rests upon the lattice and unwinds as the conveyor moves forward. Thus, asbestos covered with a layer of cotton felt is fed into the first machine of the carding unit. The carding operation has already been described.

The roving produced by the carding machines may be converted into yarn on either one of the two machines, both of which have been discussed. As was pointed out, the ring spinning machine produces a much finer yarn than the mule spinning device. Since the British specialize in fine yarn, the ring spinning frames are used virtually to the exclusion of the mule spinner. The preparation of yarn for the market and its conversion into woven products will not be discussed since they are similar to the processes used in this country.

OCCUPATIONAL ANALYSIS AND A DESCRIPTION OF THE CHIEF OCCUPATIONS

The occupational analysis of the four asbestos plants covered in the present investigation is shown in table 1.

TABLE 1.—*Occupational analysis of workers in the asbestos textile manufacturing industry*

Section and activity	Number of workers in—								Total	
	Plant A		Plant B		Plant C		Plant D			
	M	F	M	F	M	F	M	F	M	F
Preparation										
Foremen	1		2						3	
Cyclemen	1		6						7	
Weighers	3		6						9	
Feedmen	2		6						8	
Total	7		20						27	
Carding:										
Foremen	2		3						5	
Second hands	3		3						6	
Carders	9		26						30	
Strippers	3		15						18	
Tenders	3		3						6	
Total	20		45						65	
Spinning:										
Foremen	1		2						3	
Spinners	5		10						15	
Mule-spinners	5		15						20	
Ring-spinners	4		10						14	
Helpers	2								2	
Total	17		37						54	
Spooling, reeling		9		13						22
Twisting:										
Foremen	2		2						4	
Second hands	4		2				1		5	1
Twisters	15		15						30	
Winders	3		4						7	
Helpers							1		1	
Total	24		23				1		48	1

TABLE 1.—*Occupational analysis of workers in the asbestos textile manufacturing industry*—Continued

Section and activity	Number of workers in—								Total	
	Plant A		Plant B		Plant C		Plant D			
	M	F	M	F	M	F	M	F	M	F
Winding:										
Foremen	1		2						3	
Winders		13		11		1			13	25
Yarn packers	8		5						13	
Total	9	13	7	11		1			16	25
Treating: Treaters	1			2	1		5		10	2
Cloth weaving:										
Foremen	1		1				1		2	
Second hands	2		1						3	
Weavers	8		25						30	
Inspectors	1		2						3	
Calenderers	3								3	
Creelers		4		5						9
Cop winders		2								2
Helpers			1						1	
Total	15	6	30	5			1		46	11
Tape and brake-lined weaving:										
Foremen			1						1	
Weavers (tape)	6		18	4	2	1			26	5
Weavers (brake band)			2		2		9		13	
Calenderers				2	1				1	2
Creelers			7		1				8	
Clerk			1						1	
Total	6		29	6	6	1	9		50	7
Cord, rope, and braid workers			9	12					9	12
Finishing:										
Treaters and driers					1		3		4	
Calenderers					1				1	
Grinders					1				1	
Cutters and drillers					3		1		4	
Braiders					1				1	
Coilers					1				1	
Total					7		6		13	
General labor:										
Engineers	1		2		1		3		7	
Laborers	6		3				2		11	
Watchmen	2		3				1		6	
Shipping clerks			1	1	3		2		6	1
Yardmen			2						2	
Total	9		11	1	4		8		33	1
Office:										
Superintendents	1		3		1		1		6	
Others	9		8	3	3	3	2		22	6
Total	10		11	3	4	3	3		28	6
Grand total	110	28	225	56	27	5	31	1	391	90

Reference to table 1 shows that a number of men have been listed as foremen and second hands. These terms have been adopted from the cotton textile industry. They do not always mean full-time supervisory duties, as might be presumed; in most cases foremen and second hands are engaged in all the activities of the department to which they are assigned. Further mention of these occupations

PREPARATION

Crushermen empty the raw asbestos, contained in 100-pound bags, into the jaw of the crusher. They also operate the crusher and remove the asbestos fiber after crushing.

Willowers or openers handle crushed fiber and waste roving. The crushed ore is fed into the willowing machine by means of pitchforks. The willowed material is usually blown into a special bin. The waste accumulated by the machine is removed several times daily by the willower. Willowers also operate screens for separating coarse granular material (cobs) from the fiber.

Pickermen make up the asbestos-cotton batches (piling) according to specifications and pitch the material into the feeder hopper and the picker machine. The pickermen in some plants may hand truck to special bins the material which is blown from the machines.

CARDING

Carders operate the carding machines and clean them when necessary.

Stockmen handle the stock prepared by the pickermen. The material is conveyed by hand trucks from the stock bins to the card feeders.

Tenders look after the stock feeders and roving jack spools. The latter are removed when full, and replaced with empty jack spools. The occupations of tender and carder are in some plants interchangeable.

SPINNING

Mule spinners operate the mule spinning frames.

Ring spinners operate ring spinning machines.

Helpers (bobbin boys) assist the mule spinners and ring spinners conveying full jack spools of roving to the spinners and carrying empty spools back to the cards. They also transport the spun yarn and attend to the roving jacks and bobbins whenever required.

SPOOLING, TWISTING, WINDING, AND TREATING

Spoolers tend the spooling machines and handle spools on which the spun yarn is wound.

Twisters and helpers operate machines that twist spun material into heavier and stronger yarn.

Winders operate yarn winding machines.

Treaters pass yarn through special compounds (usually predominantly starch) and wind it on large reels for drying. In brake lining manufacture, treaters handle asphalt and resinous materials with which the product is impregnated.

Yarn packers wrap and box finished yarn.

CLOTH AND TAPE WEAVING

Broadcloth weavers operate both dry and wet weaving looms.

Tape weavers operate looms for weaving tape and brake lining materials.

Inspectors and calenderers inspect and examine all woven material for defects and pass tape and brake lining through calendering machines to achieve desired width and thickness.

Creelers attend to the creel racks on which the warp thread spools are mounted.

Cop winders operate cop-winding machines which wind thread on metal spindles for use in the loom shuttles.

[illegible] attend to the drying of broadcloth which has been woven

UCC 017689

A 21523

ALL OTHER

Cord, rope, and wick workers operate special cord and rope-making machines and braiding machines.

Brake-lining finishers include treaters and dryer men, calenderers who operate calendering machines for squaring uneven edges, grinders who smooth off uneven surfaces, and cutters and drillers who cut the finished band to required sizes and drill it for riveting to brake shoes for automobile and other machines. In the plants studied, these occupations were interchangeable.

General labor includes those occupations associated with maintenance of equipment and generation of steam. They also include janitors, watchmen, and shipping clerks.

Superintendence is concerned with the supervision of all plant activities.

RESULTS OF THE ENGINEERING STUDY

DUST STUDIES

In the present study, dust counts were made by the impinger method (2), using ethyl alcohol and distilled water mixtures containing from 25- to 50-percent alcohol as a collecting fluid. Two sampling methods were used for collecting dust for chemical analysis. (1) Settled dust samples were collected at the breathing level and (2) air-suspended dust was collected with a small, hand-type, vacuum cleaner fitted with a removable, paper-bag type filter. The latter method is similar to a procedure developed by Hatch (5). (See fig. 1.) In addition to chemical analyses, petrographic analyses were made on several samples. In order to determine the particle size of asbestos dust, an Owens jet apparatus was employed (2). The samples collected were measured by means of a filar micrometer at a magnification of approximately 1,000 diameters (oil immersion). Photomicrographs were also made of dust obtained by the vacuum-cleaner device described above.

NATURE OF THE DUST FOUND IN ASBESTOS TEXTILE FACTORIES

Composition.—Table 2 shows the results of chemical analysis of settled dust samples obtained by the Hatch method (5). Examination of these tables does not reveal significant differences among the various samples.[2] It will be noted, with one or two exceptions, that the total ash (no water of crystallization present) does not vary greatly. The value of 54.5 obtained for wet weaving (sample 8) is due to the moisture content of the dust sample. Calculations of composition on an ash basis bring the values into closer agreement than indicated in either table. The results given for sample 11 in table 2 are for brake lining which has been impregnated with resinous compounds.

[2] Canadian asbestos (crysotile), the asbestos used in these factories, has the following composition: Ash, 85.70; SiO_2, 40.40; Al_2O_3, 1.27; MgO, 41.41; Fe_2O_3, 2.53 percent. (Asbestos: Vol. 3, No. 7, January 1922.)

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 15 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/15

FIGURE 13.—GROSS APPEARANCE OF SUSPENDED DUST COLLECTED WITH VACUUM CLEANER DEVICE. (SEE FIGURE 1.)

A21524

Table 3.—*Chemical analyses of suspended dust samples collected in an asbestos textile plant*

Sample No.	Average chemical composition of samples (percentage)					Remarks
	SiO_2	R_2O_3[1]	CaO	MgO	Fe₂O₃	
1	35.4	9.3	1.2	31.9	6.6	General air sample.
2	35.4	9.5	1.9	36.0	6.3	Do.
3	33.9	10.1	.9	29.3	7.1	Between 2 mule spinners.
4	33.9	9.3	.7	31.6	7.0	Near ring spinning machines.
5	33.9	9.7	.7	31.8	6.7	Brookdale weaving.
6	33.9	9.7	.6	30.4	6.7	Do.
7	32.1	10.4	.4	32.1	8.3	Near twisting machines.
8	33.0	12.0	.7	33.0	8.6	Near spooling machines.
9	30.6	8.9	.7	29.6	6.8	Near winding machines.

[1] Chiefly aluminum oxide.

The losses on ignition of all samples analyzed are due to the presence of cotton fiber and moisture. The 30.5 percent loss on ignition of sample 21 in table 3 is probably due to the presence of cotton fibers. In passing, it may be pointed out that no carbonate of the order reported by Barbour and Williams was found in any of the samples analyzed (6).

No quartz was found in any of the samples examined petrographically.

The consistency of the results indicated in tables 2 and 3 is more remarkable when the physical characteristics of the dusts are taken into consideration. Figure 13 illustrates the gross appearance of the dust samples taken with the vacuum cleaner device above mentioned. Each sample shown in the figure represents two grams of dust. There are real [illegible] differences among the samples, both in color and bulk.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 16 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/16

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 17 of 57

samples, on the other hand, are bulky, the difference probably being due to the presence of long asbestos and cotton fibers in these samples. Carding and preparation samples contain cobs which are present in the raw asbestos. The darker appearance of the samples shown in the top row of figure 13 is difficult to explain. It is probably due to the




FIGURE 13.—MICROSCOPIC APPEARANCE OF SUSPENDED DUST COLLECTED WITH VACUUM CLEANER DEVICE. 200X.

contact of the spun asbestos-cotton mixture with oily metal surfaces.

The microscopic appearance of the vacuum cleaner samples is illustrated in figure 14. The presence of both cotton and asbestos fibers and of small particulate matter is readily noted. The dust from the carding room is most striking. It has few fibers and it has

UCC 017691

A21525

Particle size.—The measurement of dust particles suspended in the air of asbestos textile plants is difficult because of the presence of both asbestos and cotton fibers. The differentiation of these fibers under the microscope is not always possible, especially when the fibers are short and fine. Several samples obtained with the Owens jet apparatus were examined with a petrographic microscope in order to estimate the proportion of asbestos and cotton fibers, but no satisfactory technique could be developed for this purpose. In this study, particle size determinations based on Owens jet samples were made by three methods: (1) eyepiece projection device; (2) a filar micrometer attachment to a microscope (using oil immersion, 1,000X), and (3) by means of photomicrographs.

Data obtained with the eyepiece showing the percentage of fibers (irrespective of whether they are cotton or asbestos) are presented in table 4. These are based on Owens samples and not on impinger samples. With regard to the selectivity of the device used, little information can be given, although it is known that for discrete particles, the selectivity of the Owens apparatus and the impinger are the same (2).

TABLE 4.—*Relation of fibers to particles in various Owens jet samples*

Activity	Number of Fibers counted	Number of Particles counted	Total count	Percent fibers
Weaving [1]	83	231	314	26
Weaving [2]	45	251	296	15
Picking [2]	20	259	279	7
Carding [2]	24	254	278	9
Twisting [1]	11	237	248	4
Twisting [2]	11	206	217	5
Crushing [2]	8	192	200	4

[1] Plant B.
[2] Plant A.

The results given in table 4 indicate that weavers are exposed to the highest percentage of fibers. This is to be expected because the warp threads fray as the shuttle crosses them. Samples collected near the pickers and carders have fewer fibers, while the dusts to which crushermen are exposed have the least proportion of fibers.

The relatively low percentage of fibers in suspended dust explains the small number of fibers observed while making dust counts of impinger samples. The latter were usually diluted to give 20 to 50 particles per quarter field. Hence, except for weaver's samples in which considerable fibers were observed, fibers are only rarely found in making counts of other samples. (Dust counts included both fibrous and nonfibrous particles, irrespective of size.)

Table 5 illustrates the size range of particulate matter. Two hundred particles in each sample have been measured by means of the

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/17

Elgr nacrometer. As may be seen, the median size varies from 1.22 microns to twice this value. Fulton and his coworkers used an electrostatic precipitator to collect dust samples and calculated an average value for the total distribution of fibrous and nonfibrous of 1.33 microns (4). The median size for particles less than 5 microns, as deduced from data given by these investigators, is approximately 0.47 microns, which is much lower than the values given in table 5 of this report. This difference can probably be attributed largely to the type of sampling devices used in the two investigations.

TABLE 5.—*Size frequency distribution of particulate dust suspended in the air of asbestos textile plants*

Nature of process where sample was taken	Median size (in microns)	Geometric standard deviation	Percentage frequency of each particle size group (in microns)											
			0 to 0.49	0.5 to 0.99	1 to 1.49	1.5 to 1.99	2 to 2.49	2.5 to 2.99	3 to 3.49	3.5 to 3.99	4 to 4.49	4.5 to 4.99	5 or more	Total
Preparation	1.25	1.56	3	21	37	22	10	4	2	1	0	0	0	100
Carding	1.45	1.57	1	9	21	25	9	10	4	5	1	3	3	100
Spinning	1.30	1.24	1	1	25	34	16	4	1	5	1	1	3	100
Twisting	1.75	1.71	5	30	24	14	9	3	1	5	0	0	3	100
Weaving (broadloom)	2.30	1.91	1	4	43	22	10	3	2	5	1	1	3	100
Weaving (fine fabrics)	2.60	1.61	0	3	11	24	11	12	7	13	4	3	6	100

In determining the length of fibers present in Owens samples, photomicrographs were used. It was necessary to use this method of measurement because of the curvature and matting of fibers in the microscopic field. This difficulty was overcome by using compass pointers and measuring each fiber visible across a given section of the photomicrograph. The size distribution of fibers, classified by occupation, is presented in table 6. The fibers range from as little as 2 microns (450× magnification) to as high as 400 microns. The median sizes, as may be seen by reference to the table, range from 7 to 16.3. Fulton's data indicates (assuming all particles greater than 5 microns to be fibers) a median size of 15 microns.

TABLE 6.—*Median length of fibers sampled with an Owens jet apparatus*

Activity	Median length of fibers in microns
Willowing	7.0
Picking	9.5
Carding	8.6
Twisting	12.8
Weaving (broadloom)	16.3

No size frequency distribution with regard to the width of fibers present in the air has been made in this study. However, it has been noted, with few exceptions, that the width of fibers collected by the Owens jet apparatus was less than 0.5 micron. This is in agreement with the results given by Fulton, who shows more than 78 percent of the fibers measured by his projection method to be less than this size.

DUST CONCENTRATION IN DIFFERENT OCCUPATIONS

In earlier studies of the dusty trades carried on by the Public Health Service, determinations of the average dust concentration have been of great value (1) in describing the working environment, (2) in making comparisons of the effectiveness of different methods of dust control, and (3) in relating health conditions to dust exposure. In this study 242 impinger samples were taken and counted, and the results have been classified by occupation in tables 8 to 18. Each of the occupations carried on in asbestos textile factories has been discussed separately in the following section and data for the entire industry have been brought together in table 19. In this industry there are relatively few occupations in which workers do two or more kinds of work which expose them to totally different dust concentrations. In such cases the several exposures have been weighted according to the average time of exposure.

For example, in plant A, willowers are engaged in the following activities: willowing, 5 hours; picking, 1½ hours; storage-room work, 1½ hours. These activities are associated with exposures of 2.3, 36.3, and 16.4 millions of particles per cubic foot, respectively. Hence, we may tabulate exposures as follows and calculate a weighted average:

TABLE 7.—*Dust exposure of willower, plant A*

Activity	Time of exposure (hours per day) a	Average dust exposure (MPPCF) b	Millions of particle-hours per cubic foot a×b
Willowing	5	2.3	11.5
Picking	1½	36.3	54.4
Storage-room work	1½	16.4	24.6
Total	8		90.5

$$\frac{90.5 \text{ million particle-hours per cubic foot}}{8 \text{ hours}} = 11.3 \text{ million particles per cubic foot}$$

The average dust count of the 10 samples in the above table, calculated without regard to the duration of exposure, would be 18.7 million particles per cubic foot.

Preparation.—Preparation includes the occupations of crushermen, willowers, and pickermen. In plant B, these three occupations were conducted in the same room, while in plant A only crushers and willowers worked together. Picking (including piling), in the latter plant, was done in a separate room. The dust exposures are indicated in table 8.

UCC 017692
A 21526
Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 18 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/18

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 19 of 57
UCC 017693
A21527
asbestosclaims.law/library/18707/19
Published by AsbestosClaims.Law

Cord, rope, and braid working.—Other activities found in the asbestos textiles plants are concerned with the manufacture of special products, such as cord, rope, and braid, and in treating and finishing yarns. These operations produce little dust as may be seen by reference to table 14.

TABLE 14.—*Dust exposure of workers manufacturing and finishing special products*

Occupation	Number of samples taken	Dust concentration (MPPCF)
Cord, rope, and braid workers	8	1.2–10.2
Yarn treaters	3	1.7–3.6

The equipment used in manufacturing these products does not operate continuously. None of the operations of cord making, rope making, or braiding requires full-time personnel, and a small group of workers perform all these jobs. The highest dust count (10.2) in table 14 is for braiding. The counts obtained for yarn treaters are influenced by the operations undertaken nearby.

Brake-band finishing.—This generates little dust. In the plants studied, finishing operations were conducted in separate rooms or buildings. Dust counts are shown in table 15.

TABLE 15.—*Dust exposure of workers manufacturing finished brake-band lining*

Occupation	Number of samples taken	Dust concentration (MPPCF)	Occupation	Number of samples taken	Dust concentration (MPPCF)
[illegible]	1	1.5	Cutters and drillers	1	1.5
[illegible]	1	1.5	Inspectors	1	1.5
[illegible]	1	1.2	Coilers	1	1.5

Workers employed in the occupations listed in this table are exposed to dust which differs in composition from the dust found elsewhere in asbestos textile plants. This is because the dust generated by grinders, cutters, and drillers of treated stock is made up of resin, asphaltum, asbestos, and metal. (See analytical results for sample 11, table 2.)

Miscellaneous occupations.—In addition to the groups directly concerned with the manufacture of textiles, a considerable number of workers (engineers, general laborers, clerks, and superintendents) are engaged in other activities. With the exception of certain foremen, these workers are exposed to low dust concentrations. The data presented in table 16 require no comment.

TABLE 16.—*Dust exposure of miscellaneous groups*

Occupation	Number of samples taken	Dust concentration (MPPCF)	Occupation	Number of samples taken	Dust concentration (MPPCF)
Engineers	2	1.2	Yardmen	2	0.10
Laborers	2	.36	Foremen (plant)	(1)	(1)
Watchmen	1	1.2	Superintendents (other)	2	.12
Shipping clerks	1	.12	Clerks	2	.12

(1) Samples and exposures depend on the activities with which they are associated.

PREVIOUS DUST EXPOSURE

The dust concentrations discussed in the preceding paragraphs represent present-day conditions. The officials of asbestos textile plants have in recent years attempted to control the dust hazard. In the plants studied, efforts have been made to exhaust picking, carding, and brake-band grinding machines. (Another plant, which will be discussed later, has extended the use of control equipment to nearly every dusty operation concerned with the manufacture of asbestos textiles.) A knowledge of the exposure of workers prior to the introduction of control equipment is important in correlating clinical findings with dust exposure.

In order to estimate past exposure, the air flow through various hoods attached to carding machines in plant A was shut off for 2 hours. Dust was allowed to accumulate for 1 hour prior to sampling. The dust thus generated caused much inconvenience to employees in the course of their work. For this reason, permission to shut off exhaust equipment was not granted in other plants studied. Carding machines are sensitive to slight air movements and are consequently difficult to regulate. Once they have been adjusted, any external disturbance affects the weight of the roving produced.

The dust counts obtained under conditions simulating those prior to the installation of the exhaust system are shown in table 17. For contrast, present exposures are included.

TABLE 17.—*Dust counts when control measures are used and when they are not in use*

Occupation	Average dust count—past (MPPCF)	Dust concentration—present (MPPCF)	Occupation	Average dust count—past (MPPCF)	Dust concentration—present (MPPCF)
Crushermen	50.2	2.3	Mule spinners	15.5	2
Willowers	50.2	11.1	Ring spinners, twisters, spoolers, and winders	3.9	1.2
Pickermen	83.0	31.3	Weavers	7.1	1.7
Carders	72.3	29.1			

The low counts obtained for the occupations listed below carders is due to a partition which separates them from the carding machines.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 20 of 57

Mule spinners are located immediately beyond this partition, and the dust count shown is due to dust which has passed through an open door.

Three dust samples obtained in the carding room 1 hour after the exhaust fans were shut off gave counts of 77, 73, and 78 million particles per cubic foot. At the end of 2 hours a sample taken in the same room gave a count of 61 million particles.

No major changes have been made in plants C and D since the start of operations. The dust counts made are essentially the same as those which have always existed. On the other hand, no data are available for past exposures in plant B. This failure can be attributed to several factors, but principally to the gradual expansion of the plant with the growth of the industry, which increased the amount of equipment, and the changes made in recent years with the exhaust system. The first of these factors appears to have affected all operations conducted in the plant, while the latter is restricted to the carding and preparation departments. In addition, it may be pointed out that plant expansion has tended to increase dust productivity, while the introduction of exhaust systems has reduced it. Hence, it may be assumed that the exposures of all occupations in plant B following carding were probably higher during the study than prior to expansion. It must also be remembered that more men have been working since the introduction of added equipment than at previous times.

UCC 017694

INTERPRETATION OF DUST COUNTS

The average dust concentrations just discussed make a satisfactory measure of dust exposure for most purposes. In correlating medical findings with dust exposure, however, it is frequently advantageous to express the dust exposure in units which combine duration of exposure with average dust concentration. An easily calculated descriptive value was developed during statistical analysis of the results of an engineering and medical study of anthraco-silicosis (51). If a worker's dust exposure has been unchanged for all practical purposes during his period of employment the average dust count (in million particles per cubic foot) is multiplied by the duration of exposure in years. The product is used as a numerical measure of his total dust exposure. If a worker has changed his occupation (and in so doing has changed his dust exposure appreciably) a calculation such as the one shown in table 18 is made.

A21528

TABLE 18.—*Example showing the method of computing dust exposure of cluster textile workers*[1]

Occupation of worker	Number of years in occupation	Dust count (MPPCF)	Millions of particle-years per cubic foot of air
Carder[2]	3	72.3	216.9
Carder	2	29.1	58.2
Mule spinner	2	2.6	5.2
Total	7		280.3

[1] Plant A.
[2] Prior to installation of exhaust system.

Weighted dust exposure = $\frac{280.3 \text{ millions of particle-years per cubic foot}}{7 \text{ years}}$ = 40 million particles per cubic foot

The worker whose dust exposure has been calculated in table 18 would be assigned a value of 280 million particle-years in tabulations in which, for the sake of simplicity, it was desirable to have a one-term measure of dust exposure; table 42, for example. When average dust concentration and duration of exposure are to be tabulated separately, as in table 43, the individual described in table 18 would be assigned a dust concentration of 40 million particles per cubic foot.

THE CONTROL OF ASBESTOS DUST

The control of dust in industry often entails many practical difficulties. These difficulties pertain chiefly to the handicaps imposed by control equipment upon the performance of the work being done, and in estimating a priori the air volumes required. Both of these difficulties are inherent in the asbestos industry. The arrangement necessary for removing dust from carding machines and from looms, which demand accessibility to all parts, is self-evident. Added to this requirement is the limited information available with regard to suitable design of equipment for each dust-producing source and the requisite air volumes. The latter item is most important in the design of hoods for carding machines, inasmuch as air velocities must be kept relatively low in order to prevent the asbestos-cotton mixtures from being subjected to air currents which convey them to the exhaust system. At the same time, however, the velocities chosen must be adequate to maintain the dust below the safe or threshold limit.

The dust-control data contained in the following paragraphs are based on observations made in two plants included in the present study and on data obtained elsewhere by Assistant Public Health Engineer Richard Page (7). The data presented include operations for which control methods have been developed. In addition, the type of control equipment used in England is discussed, although no values pertaining to air requirements are available in this instance.

PREPARATION DEPARTMENT

The dust counts presented earlier in this bulletin for the preparation department apply to uncontrolled conditions. A very high degree

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/20

...spread can, however, be effected under certain conditions, especially by the use of willower hopenings, piling operations, and picking. The crushing machines in the plants studied were not exhausted or segregated. In England, on the other hand, it is customary to enclose and segregate crushing equipment while it is in operation.

Willowing operations.—Willowing machines are often productive of high dust concentrations (36 MPPCF). As has been previously mentioned, this dust is caused both by the manual methods used in feeding these machines and by the action of the machines themselves. Consequently, a reduction of dust concentration may be obtained by using a conveyor feeder which eliminates raising materials to waist




FIGURE 15.—EXHAUSTED PILING BINS.

levels with forks and using an exhaust above the feeder box and below the machine housing. The willowed material should, of course, be pneumatically conveyed to a suitable bin which is segregated from the preparation room.

In the case of two machines equipped with exhaust, it was found possible to reduce dust concentrations to 3.6 MPPCF. The exhaust hoods located above the feeders drew 400 CFM, while the downward exhaust below the willower housing drew amounts of air ranging from 225 CFM to 600 CFM, the latter air flow being used on a fly willower. (See fig. 4.) The air handled by the pneumatic conveyor was 1,800 CFM. For the 11-inch diameter ducts used, this corresponds to an air velocity of 2,750 feet per minute.

Piling.—Piling operations are usually conducted close to picker machines. The dust generated by this operation may be effectively controlled by using small exhausted compartments or bins. Compartments constructed as shown in figure 15 and handling approximately 1,025 CFM of air reduce the dust concentration from an average of 20 to 5.4 MPPCF.

English practice in piling consists in using exhaust manifolds close to the mixture being made (fig. 16). Air volumes are not available for the type of exhaust shown in the figure. The method appears to be cumbersome and to interfere with the process being carried on.

Picking.—The exhaust control used on picker machines is partly




FIGURE 16.—EXHAUST VENTILATOR USED FOR PILING OPERATIONS (BRITISH PRACTICE).

shown in figure 17. One hood is connected at the feeder end and one at the discharge end. The center portion of the picker is also exhausted (not shown in figure). With this type of arrangement the dust concentration averages 6.7 MPPCF. This value is considerably lower than found among uncontrolled equipment which exceeds in many instances 74 MPPCF.

It was found that hoods above the feeder handled 500 CFM; the downward exhaust at the center measured 1,620 CFM; at the discharge end the air flow was 450 CFM. Pneumatic conveyors attached to the discharge end of some of the pickers handled 2,000 CFM. These ducts were approximately 12 inches in diameter, thus providing for a conveying velocity of 2,550 feet per minute.

UCC 017695

4215 29

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 21 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/21

[illegible] special conveyor belts operating between the [illegible] the feeder mixer that the dust concentrations might be reduced below the value of 6.4 MPPCF, indicated above.




FIGURE 17.—SCHEMATIC DRAWING OF EXHAUST SYSTEM USED WITH PICKER [illegible]

CARDING

Two types of dust-control equipment are at present used on carding machines. The first type consists of a hood which is located directly above the exposed brushes of the machine (fig. 18), while the second type is totally enclosed and exhausted at various points (fig. 5). Studies made of the first type of exhaust systems showed a reduction in dust concentration ranging from 72.3 MPPCF when no ventilation was applied to 29.1 with ventilation. The air flow for each hood averaged 870 CFM, which corresponded to six air changes per hour in the room in which the cards were located. It may be seen that the type of hood and the air flow used reduce the dust concentration to approximately one-third the amount created without ventilation.




FIGURE 18.—VERTICAL EXHAUST VENTILATION APPLIED TO AN UNENCLOSED PRIMARY CARDING MACHINE.

The second type of exhaust system used proved more effective than the first in reducing dustiness. The dust concentration determined from numerous samples ranged from 1 to 3.9 million particles per cubic foot. This high degree of dust removal was accomplished with an arrangement similar to that indicated in figures 19 and 20. In the primary [illegible] *a* and *b* discharge upward. The duct *c* exhausts from the near the center of the machine. The openings *e*, *f* and *f'* are located at the bottom of the cards and exhaust the bottom fly which accumulates. The air volumes exhausted by each opening are indicated in the figure.

In addition to careful enclosure of the machine proper in the second type of system used, the feed box is carefully covered.




FIGURE 19.—TOTALLY ENCLOSED EXHAUSTED PRIMARY CARDING MACHINE.

The design of an exhaust system for carding machines is rendered exceedingly difficult owing to the ease with which carding material leaves the rolls. High air flows may seriously affect the quality of the roving produced. Likewise, a poorly constructed housing may create eddy currents which may prove bothersome. Any system developed requires care and experiment in design.






FIGURE 20.—ELEVATION AND PLAN FOR SPOOLING EXHAUST SYSTEM.

SPOOLING AND TWISTING

The dust generated by spooling operations may be [illegible] of cones in which the spools are inserted and exhausted. Such a system is shown in figure 21. Each case is constructed so as to offer a minimum amount of interference to the work being done. In the case of four Foster winders (200 spools) each cone is 9 inches in diameter across its widest part and 2 inches at the duct connection. The cones are each 10 inches deep. Two hundred cones were found to handle an

UCC 017696


A21539

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 22 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/22

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 23 of 57

...270 CFM or an average of 46.5 CFM per cone. The ... operation was 2.9 MPPCF. Without the cone system ... above, the dust concentration for the same operation was ... MPPCF.

Twisting machines are more difficult to exhaust locally than spoolers. Figure 22 shows a system which has been partly completed. It consists of exhausting the lower section of the twisting machine by extending a duct with openings along the entire length of the equipment. Each system attached to the machine is intended to



UCC 017697

FIGURE 21.—EXHAUSTED SPOOLING FRAME.

exhaust 1,670 CFM. Dust determinations to establish the effectiveness of this system could not be made owing to the fact that only one of 10 twisting machines was equipped with exhaust and the counts obtained were affected by unexhausted machines operating close by.

WEAVING

Any system developed for removing dust generated by broad looms represents the accomplishment of one of the most difficult operations encountered in the design of exhaust equipment. Nevertheless, this has been done effectively both in this country and in England. The American system shown schematically in figures 23 and 24 consists of a downward-draft hood fitting close to the shuttle rack of the loom and a movable upper exhaust hood which operates on the surface of the woven material. The latter hood is actuated to and fro by the action of the loom lay to which it is attached. The hood is connected to the main exhaust duct by means of a flexible air tight joint. Both the downward and upward exhaust hoods are narrow and possess the advantage of high opening velocities. The upward-draft hood connected to the loom lay is segmented. Each hood used, as indicated in figure 24, handles 1,300 cubic feet of air per minute and reduces the dust concentration to an average of 0.7 MPPCF. The system is thus extremely effective, since on uncontrolled looms the concentration averages 49.7 and may exceed 140 MPPCF.




FIGURE 22.—EXHAUSTED TWISTING FRAME.

The English method for exhausting looms differs considerably from American practice. In the former system all parts are stationary (fig. 25). The looms are exhausted upward by hoods located in front of the operator and resting a short distance away from the lay. The hood consists of two parts which may be turned back whenever repairs to the loom are necessary. No data as regards air volumes used in this type of equipment or its effectiveness in reducing dust are available.

INSPECTION

In some plants, asbestos cloths are brushed during inspection. The usual procedure consists in brushing the cloth from above and below as it passes over the inspection table. An inspection table of this type with exhaust hoods is shown in figure 26. The hoods used are located near the cloth brushes, and extends the full width of the cloth examined. They are, however, only one-half inch in width.

A study made of an inspection table such as discussed above indicated that two hoods located on either side of the cloth handled approximately 730 CFM each. A third brush and hood located near

A21531

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/23

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 24 of 57

...ed at the end of the inspection table handled 192 CFM. The ... reduced the dust generated from a value of 11.1 when no ... was applied to a value of 0.5 MPPCF with ventilation.

SUMMARY OF THE STUDY OF THE WORKING ENVIRONMENT

The engineering data contained in this report are based chiefly on a study of four asbestos plants. Data on dust control methods



FIGURE 23.—EXHAUSTED BROADCLOTH LOOM.

have been secured in the course of a separate study undertaken in a fifth plant. Two of the four plants investigated were engaged in the manufacture of asbestos textiles directly from raw crysotile (Canadian) asbestos received in 100-pound bags. The remaining two plants began operations with yarn received from other sources and were concerned only with the manufacture of asbestos, tape, braid,

UCC 017698

A21532

Dust concentrations were determined by means of the impinger technique using light field counting. In all, 212 impinger samples were taken so as to evaluate the environment of each occupation.




FIGURE 24.—DETAIL OF EXHAUSTED LOOM SHOWN IN FIGURE 23.




FIGURE 25.—DETAIL OF EXHAUST DUCT ON BRITTEN TYPE LOOM.

The number of workers employed, the number of dust samples taken, and the dust concentrations determined for each occupation are presented in table 18. The concentrations range from values of 0.86




FIGURE 26.—EXHAUSTED BRUSHING AND INSPECTION TABLE.

for shippers to as high as 74.3 MPPCF for pickermen. Occasional samples exceeding 100 MPPCF were found for the occupations of carding and weaving.

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/24

A21533

UCC 017699

2. Since material leaving picking machines is productive of considerable dust, storage rooms receiving such material should be exhausted. Workers who must enter such rooms while material is being blown should be equipped with approved types of respirators. No material should be removed from storage rooms while the picking machine is in operation.

3. Clean air should replace the dust-laden air removed by exhaust systems attached to the equipment. This requires, in event of recirculation of air, that dust concentrations entering any room be kept at levels comparable to that found in outdoor air.

4. Periodic studies of the condition of the working environment appear necessary to determine whether the control methods adopted are constantly adequate. This requires that dust concentrations be determined for each operation. Ventilation measurements are equally important. The work of inspection and review should be performed by persons trained for such studies.

UCC 017700

A21534

STATISTICAL DESCRIPTION OF EMPLOYEES

OCCUPATIONAL DISTORIES

At the start of the medical examination the physicians questioned each worker to obtain a complete record of all the jobs at which he had worked. This information was recorded on the portion of the physical examination form reproduced in figure 27. After the basic census data (name, age, sex, color, and marital status) had been recorded the worker was asked how old he was when he first began

CASE RECORD

Name ..... Date of Birth .....
(Surname) (First Name)

Address .....

Age ..... Sex ..... Color ..... Racial Stock ..... M. S. W. F. Sp.

Plant ..... Address .....

Age began work ..... Years worked ..... Time idle .....

OCCUPATIONAL HISTORY

SPECIFIC JOB	INDUSTRY	YEARS		WHERE	WORKING CONDITIONS AND CONTROL MEASURES USED
		FROM	TO		
Pres.					
Prev.					
"					
"					
"					
"					
"					
"					
"					
"					
"					
"					
"					
Total Years					

FIGURE 27.--FORM USED FOR RECORDING OCCUPATIONAL HISTORY

industrial work. This figure was subtracted from his present age, and a line of questioning was begun to elicit information on the work done in the intervening years. No occupational history was regarded as complete until all of this time could be accounted for by definite entries. The job he held at the time of the study was entered on the top line of the occupational history, together with the number of years he had held it, the name of the department in which he worked, and any other pertinent data that could be obtained. Next, he was asked what he had done previously, and again a complete entry was made. The name of the factory was entered for each period of em-

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 26 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/26

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 27 of 57

UCC 017701

[illegible] industry [illegible] After each job had been [illegible] described what he had done before that. [illegible] had worked in [illegible] industry in which workers are known to be exposed to pneumoconiosis-producing dust, entries were made in the last column to show what materials he handled and whether or not dust-control measures were in use. Information on previous employment in dusty trades is of paramount importance in a study such as this. Any cases of prolonged exposure to pneumoconiosis-producing dust other than asbestos must be treated separately.

After the field study had been completed, an engineer assigned dust exposure values to each worker, following a procedure illustrated in table 18. In assigning these values, consideration was given to each job a worker had held in an asbestos plant.

The importance of the occupational history in a study of this kind can scarcely be overemphasized. It is the connecting link between the engineering and medical studies. The correlations between medical findings and dust exposure that adequate occupational histories make possible provide means for estimating the dust concentrations to which workers may be exposed safely.

Only 23 persons reported previous employment in an industry in which workers are known to be exposed to pneumoconiosis-producing dust. Such periods of employment usually were so short that no deleterious effects would be expected. Twenty-one persons had normal lung-field markings. Only one of these 23 persons had asbestosis at the time of this study, a man who had worked 3 years in a foundry and 10 years in asbestos textile plants.

COMPARISONS WITH OTHER INDUSTRIAL WORKERS

Five-sixths of the employees were native-born Americans of Anglo-Saxon stock, and the remainder were Negro males.

Before beginning work in asbestos textile factories, more than 200 of the 541 persons had worked at similar occupations in cotton or wool textile plants. A large proportion of the employees had worked on farms either before beginning industrial work or in intervals between periods of industrial employment.

About 15 months before this study was begun, approximately 150 workers in these asbestos textile factories were replaced by workers with little or no previous asbestos exposure. Although an effort was made to examine as many as possible of these former workers, less than half could be examined. Partly as a result of these personnel changes, the group of employees examined in this study had an unusually low average age. There was an abnormally large percentage of workers with less than 5 years' employment in the asbestos textile industry and an abnormally small percentage of persons who had worked 10 years or more in this industry.

TABLE 21.—*Age distribution of* [illegible]

Age	Number of workers			Total
	White males	White females	Negro males	
Total	338	118	85	541
Under 20 years	10	12	7	29
20 to 24 years	53	26	17	96
25 to 29 years	57	28	23	108
30 to 34 years	51	18	12	81
35 to 39 years	54	13	13	80
40 to 44 years	36	7	3	46
45 to 49 years	44	5	4	53
50 to 54 years	7	6	4	17
55 to 59 years	10	3	1	14
60 to 64 years	2	0	0	2
65 to 69 years	1	0	0	1
70 years and over	0	0	0	0
Unknown age	1	0	0	1
Average	42.08	39.36	42.41	[illegible]
Standard deviation	9.39	10.19	8.85	[illegible]

The low average age of this group of asbestos textile workers must be kept in mind in making comparisons of health conditions in these factories with health conditions in American industry in general.

TABLE 22.—*Average age of white male industrial workers*

Industry	Average age	Public Health Service Bulletin	Industry	Average age	Public Health Service Bulletin
Asbestos textile	42.1	[illegible]	Glass	42.3	[illegible]
Iron and steel	42.0	202	Gas	41.8	[illegible]
Granite cutting	39.9	187	Coal	41.2	[illegible]
Anthracite mines	39.5	221	Cement	41.0	[illegible]
Fur cutting	39.2	234	Rubber	40.1	[illegible]
Pottery	38.1	136	Chemical	39.1	[illegible]
Post office	37.8	162	Garment	32.8	[illegible]
Foundry	37.8	162			

The average age of white male industrial workers is less in the asbestos textile factories studied than it is in 14 other industries in which the Public Health Service has made similar studies. White males employed in asbestos textile factories average 32.1 years, white females 30.4 years, and Negro males 32.4 years of age.

TABLE 23.—*Length of employment in the asbestos textile industry*

Sex and color	Total	Years employed in asbestos textile industry						
		Less than 1	1 to 4.9	5 to 9.9	10 to 14.9	15 to 19.9	20 to 24.9	25 to 29.9
Total	541	73	260	102	47	16	2	1
White males	329	41	164	72	28	13	2	1
White females	111	19	53	19	13	4	0	0
Negro males	79	13	43	11	6	2	0	0

A21535

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/27

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 28 of 57
UCC 017702
A21538
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/28

...pneumoconiosis, the average length of employment in the asbestos ... industry is relatively short. Only 18.8 percent of the employees ... been employed in the asbestos textile industry more than 10 ... In the 13 industries included in table 22 for which information ... length of employment is readily available, only two industries have ... a percentage of employees who have worked less than 10 years. ... is partly accounted for by the fact that the asbestos textile ... are of comparatively recent origin in this country. Only one ... plants studied had been in operation before 1920.

MEDICAL STUDY

PROCEDURE

A medical examination was made of each worker employed in the ... asbestos textile factories of North Carolina during the winter of ... 1930. A group of former asbestos workers was also examined in ... order to obtain a complete sample of asbestos-exposed workers. ... workers and other employees not exposed to asbestos dust were ... studied in these examinations. They are classified as controls. ... included in this control group, which numbered 76 persons, were 27 ... persons seeking employment in the asbestos industry for the first ...

Before making any medical examinations the two physicians made ... inspection tour of each factory in which this study was made in order to familiarize themselves with the various jobs or occupations involved in the manufacturing process. This is a necessary prerequisite for taking a usable occupational history.

... space for examination and fluoroscopic rooms and laboratory was obtained at a ... as convenient as possible for all concerned. An improvised darkroom was ... for loading and for processing the X-ray films. In an adjoining room, the ... medical examinations were made by two physicians approximately according to ... following schedule: Four persons at 9 a. m.; four at 10:30 a. m.; four at 1 p. m.; ... at 2 p. m.; and four at 3 p. m. In other words, about 30 minutes were required per examination. The routine used in the study of pulmonary tuberculosis ... followed in the examination of the chest. After the history and physical ... had been recorded, the group of four was fluoroscoped and chest roentgenograms taken. A few examinations were made in the evening, and other ... were made in the schedule so as not to interfere too seriously with the ... routine of the employees.

At plant A, the urine of workers was examined with particular attention to ... of silica excretion. In addition to this metabolic study, routine examinations were made of the sputum of workers in plants B and C for presence of acid-fast bacilli and asbestosis bodies.

Sections of the physical examination form used in this study have been reproduced in several places in this bulletin as a means of indicating the information that was collected.

PAST MEDICAL HISTORY

Each person was questioned in order to find out whether he had previously had certain diseases which are of importance in the study of chest diseases.

FAMILY AND PAST MEDICAL HISTORY (Insert Date when Positive)

Family history of T. B.....

Exposure to T. B.....

Tuberculosis..... Treatment..... Pleurisy..... Dry..... Wet..... Site.....

Hemoptysis..... Dates..... Amount..... Asthma..... Hay Fever.....

Pneumonia..... Lumpsome..... Bronchitis..... Sinus Disease.....

Influenza..... In Bed..... More colds in recent years.....

Average number of colds yearly..... Head ..... Duration..... Severity.....
Chest .....

Tonsilitis .....Previous injuries .....

Operations .....Heart Disease .....

Rheumatic Fever..... Scarlet Fever..... Hypertension.....

G. I. .....Skin disease .....

G. U. .....Nervous system .....

Menstrual .....Pregnancies .....

Other serious illnesses or complaints.....

.....

PRESENT HEALTH

Cough ..... How long.....Type..... When worst.....

Productive..... Amount.....Character..... Odor.....

Dyspnea..... Duration.....Severity..... Progressing.....

Appetite..... Digestion.....Bowels..... Edema.....

Strength and energy..... Nerves.....

Weight: Stationary..... Lost..... Gained ..... How long.....

Other complaints .....

.....

Progress of symptoms .....

.....

.....

HABITS

Alcohol .....Sleeping .....

Tobacco .....Eating .....

FIGURE 20.—FORM USED FOR RECORDING PAST AND PRESENT MEDICAL HISTORY.

Tabulations of previous illness did not indicate that pneumonia, pleurisy, respiratory diseases other than the common cold, digestive

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 29 of 57
UCC 017703
A21537
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/29

a fortnight or prolonged contact with an active case of tuberculosis and made people more susceptible to asbestosis. Frequent, disabling colds were more common among persons with advanced pulmonary fibrosis than they were in persons with little or no fibrosis, but this probably is a result of the fibrosis rather than a predisposing influence.

SYMPTOMS OF ASBESTOSIS

The chief symptoms usually associated with asbestosis are progressive dyspnoea, cough, hemoptysis, emaciation, weakness, poor chest expansion, curved finger nails or clubbed finger tips, chest pain, and cyanosis. In these respects asbestosis resembles silicosis. Early cases, of course, would have relatively few of these symptoms and an advanced case would have almost every symptom listed. The appearance of these symptoms can be explained on the basis of a more or less general fibrosis of the lungs and concomitant pathological changes such as emphysema, bronchiectasis, and fibrous pleurisy. The cardinal symptoms of asbestosis are cough and dyspnoea. These two symptoms occurred together and in combination with other symptoms far more frequently than one would expect on the basis of chance alone.

TABLE 24.—*Number of persons who had certain pairs of disorders, compared with the number who would be expected to have them on the basis of chance alone. Where the differences are statistically significant the observed numbers appear in bold-face type*

Disorder	Persons affected		Cough		Hemoptysis		Loss of weight		Chest pain		Weakness		Loss of appetite	
	Number	Percent of total in group affected	Observed	Calculated[1]	Observed	Calculated	Observed	Calculated	Observed	Calculated	Observed	Calculated	Observed	Calculated
Dyspnoea	122	9.3	91	75.6	**30**	19.2	25	17.2	11	5.0	31	21.1	44	15.6
Cough	1,312	100.0			**32**	11.6	**9**	12.2	23	5.7	18	12.4	34	13.5
Hemoptysis	32	2.4							11	1.3	9	3.7	9	4.1
Loss of weight	25	1.9												
Chest pain	23	1.8												
Weakness	49	3.7												
Loss of appetite	44	3.4												

As table 24 shows, 91 of the 122 people who complained of dyspnoea had a cough also. On the basis of chance alone, assuming there was no causal connection between the two symptoms, one would expect that 36 people would have both conditions.[3] Actually, the two conditions are found together far more frequently than that, and it is safe to assume that there is some sort of causal relationship. Similar considerations hold for most of the other pairs of impairments.

Typically, the cough of asbestosis is dry and nonproductive. In some cases there is scanty, viscid expectoration. Severe paroxysms of coughing occasionally occur in cases of asbestosis having serious disability. At such times highly tenacious sputum is raised.

Usually a complaint of dyspnoea was entered on the records only when breathing was labored after an exercise test (see p. 65) carried on in the presence of the examining physicians. The high percentage of persons with dyspnoea who were able to be at work is undoubtedly due to the sedentary nature of much of the work in asbestos textile factories.

Blood-streaked sputum occurred frequently among persons exposed to asbestos dust although there were only two active cases of pulmonary tuberculosis in the group. These two cases have not been included in tables 25 and 26 and figures 29 and 30. The absence of pulmonary tuberculosis makes hemoptysis a more significant finding. Its presence seems to indicate that bronchiectatic and emphysematous processes are going on in the lungs. The presence of asbestosis bodies in the sputum (which will be discussed in a later section) also can be interpreted as a sign of disintegrative processes in pulmonary tissue.

Weakness and loss of weight are closely related symptoms. Independently of age and other factors, these symptoms increase in frequency with increasing dust exposure. They are accompanied by a general bodily appearance not unlike that of fibroid phthisis.

Nineteen of the 447 asbestos-exposed people complained of a pain in their chests which is best described by the term "substernal distress." It was vaguely defined and not as a rule very well localized, and it seemed to be an ache or a feeling of oppression rather than a sharp pain. It did not seem to be of cardiac origin. Only one of the 24 persons with a past history of pleurisy complained of a chest pain of this kind. It appears that substernal distress may be connected with the cough or the dyspnoea that accompany asbestosis (table 24).

The symptoms of asbestosis are listed in table 25, where their incidence is classified according to the total dust exposure of the persons affected. For purposes of comparison, the employees of asbestos textile factories have been divided into five groups, which differed in dust exposure, one of them being a control group of employees who were not exposed to asbestos dust. A number representing the total dust exposure was assigned to each dust-exposed employee by multiplying the number of years he had worked in the asbestos industry by the average dust concentration to which he presumably had been exposed. His total dust exposure is thus expressed in million particle years. If a worker had changed from one occupation to another, a dust exposure value in million particle years was calculated for each occupation, and these values were added together to obtain his total dust exposure.[4]

[1] This value has been calculated on the assumption that 0.251 represents the probability that any one of the 1,312 persons would have dyspnoea and 0.7 of 1 percent represents the probability that a person would have a cough. The 1,312 persons are taken from the second column of figures of table 24. Then, 0.251 × 0.2981 × 1,312 = 76.35.

[4] An example (table 19) has been given in the section entitled "Interpretation of dust counts."

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 30 of 57

UCC 017704

The percentage of workers who complain of cough, dyspnoea, blood-streaked sputum, and other symptoms increases rapidly with increasing asbestosis-dust exposure. In all cases the percentage of workers who are affected by a particular symptom is larger than the percentage in the control group. Two of the cases of dyspnoea found




FIGURE 29.—PERCENTAGE OF PERSONS, CLASSIFIED BY DUST EXPOSURE, WHO HAD CERTAIN DISORDERS.

in the control group were accompanied by heart defects severe enough to have caused shortness of breath.

TABLE 27.—*Classification of present complaints according to the dust exposure of the affected persons*

Complaint	Not exposed to dust		Million particle years dust exposure								Total
			Less than 25		25 to 49		50 to 99		Over 100		
	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
Total	50	100	198	100	85	100	55	100	105	100	593
Cough	6	8	24	11	23	27	19	35	61	61	110
Dyspnoea	5	7	24	12	23	27	14	53	62	59	122
Blood-streaked sputum	0	0	10	5	7	9	5	9	18	18	40
Loss of weight	4	5	11	6	8	9	3	5	16	15	38
Weakness	4	5	11	6	8	9	5	9	22	21	46
Heart defect or dyspnoea	0	0	3	2	3	4	2	4	16	15	24
Chest pain	1	1	6	3	4	5	3	5	11	10	25
Loss of appetite	0	0	0	0	2	2	2	4	2	2	6

[1] Data not available for all workers.

Another indication that these symptoms are an integral part of [illegible] is found in the relation of these symptoms to the fibrotic changes that occur in the lungs. The degree of fibrosis can be estimated from the lung-field markings seen on X-ray examination, a subject which will be more fully discussed in a later section. For the present it will be sufficient to say that as a result of long-continued exposure to asbestos dust the normal or first-degree exaggeration of linear pulmonic markings become coarser and more widespread (second-degree linear exaggeration). Upon longer or more intensive exposure a fine, interstitial fibrosis sets in which gives a hazy, ground-glass appearance to the lung field (first-




FIGURE 30.—PERCENTAGE OF PERSONS, CLASSIFIED BY LUNG-FIELD MARKINGS, WHO HAD CERTAIN DISORDERS.


degree ground glass) and eventually almost completely obscures the linear markings (second-degree ground glass).

Asbestos-exposed people have been classified into four groups according to their lung-field markings, and the percentage of persons having certain symptoms has been calculated for each group. Here again, the incidence of these symptoms increases greatly with increasing fibrotic change. Such a relation is entirely to be expected for dyspnoea, since the disease diminishes the volume of the lung available for the respiratory function. The high incidence of the other symptoms in the presence of fibrotic changes that have progressed to the ground-glass stage is additional evidence of degenerative changes in the lungs.

A21538

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/30

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 31 of 57



...according to lung-field markings of ...

Percent composition	First-degree disease		Second-degree disease		First-degree round films		Second-degree round films		Total
	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
Total	32	20	20	12	80	49	22	13	162
	32	22	20	14	36	25	55	38	143
	0	0	0	0	44	73	16	27	60
	0	0	0	0	11	27	30	73	41
	0	0	0	0	3	7	37	93	40
	0	0	0	0	25	68	12	32	37
	0	0	0	0	10	100	0	0	10
	0	0	0	0	0	0	0	0	2

X-RAY FINDINGS

LUNG-FIELD MARKINGS

The classification of lung-field markings used in the preceding section is part of a scheme that was used in a study of anthracosilicosis

FLUOROSCOPIC EXAMINATIONS

Examined by: .....

X-RAY INTERPRETATION

Technic ..... chest cm.

Examined by: .....

UCC 017705

A21539

FIGURE 31.—FORM USED FOR RECORDING X-RAY FINDINGS.

(32, 34). It is, of course, highly desirable to have a system of classification which will help in deciding whether a given type of lung-field marking represents an early or a late stage of a pneumoconiosis, and this scheme has been drawn up to illustrate the probable sequence of lung-field markings that occurs during long-continued exposure to a pneumoconiosis-producing dust. During medical studies of anthracite miners, for instance, all of the stages represented in figure 32 were found. Among asbestos-exposed workers, on the contrary, no case was found which had progressed beyond the ground-glass stage. This constitutes a difference between asbestosis and silicosis that Merewether (25), Lanza (18), and Pancoast (29), among others, have commented on. Further investigation will be needed to decide whether this difference is due to a peculiarity of asbestos dust or




FIGURE 32.—SCHEME USED IN CLASSIFYING X-RAY MARKINGS

whether it is due to the rare occurrence in asbestos textile mills of the excessively high dust exposures (100 to 300 million particles per cubic foot and more) found in certain other dusty trades.

In a typical case of asbestosis the lower half of the chest X-ray film has a fine, granular appearance which Merewether (25) aptly describes as "ground glass." Not infrequently, there is more fibrosis on the right side than on the left. The presence of emphysema in the apices, which results in increased radiolucence, tends to exaggerate the cloudiness of the bases of the lungs. Emphysema precedes the ground-glass appearance of the bases. Chest X-rays of individuals with pneumoconioses induced by other silicates, with the exception of mica,[5] do not regularly show the basal localization of fibrosis that is so characteristic of asbestosis.

[5] Unpublished data.

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/31

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 32 of 57
UCC 017706
A21540
asbestosclaims.law/library/18707/32
Published by AsbestosClaims.Law

Occasionally, in an asbestotic chest, the heart has a shaggy outline when viewed roentgenographically, a phenomenon discussed by other investigators. This seems to be of more frequent occurrence when the worker has been exposed to dust containing a high proportion of fibers (the dust produced in twisting or broadcloth weaving, for instance) than when he has been exposed to short asbestos particles,




FIGURE 31.—PERCENTAGE OF MALES, CLASSIFIED BY AGE, WHO HAD CERTAIN LUNG-FIELD MARKINGS. 229 MEN HAD NO PREVIOUS INDUSTRIAL DUST EXPOSURE, 80 HAD BEEN EXPOSED TO KAOLIN DUST, AND 357 HAD BEEN EXPOSED TO ASBESTOS DUST

such as those produced in carding. Further study would be necessary in order to establish this point.

Unlike silicosis, nodular and conglomerate shadows are not found in asbestotic workers in the absence of tuberculosis or other infections. The principal difference discernible on an X-ray film between an early and an advanced stage of asbestosis lies in the density of the ground-glass markings and the degree to which the linear pulmonic markings have been obliterated. Although first-degree and second-degree ground-glass markings have been tabulated separately in several particular film represents an early or an advanced stage of pulmonary fibrosis.

There is, of course, an increase in the prevalence of lung fibrosis with advancing age. Because ground-glass appearance of the lung fields, which is indicative of fine, interstitial pulmonary fibrosis, is a most important diagnostic sign of asbestosis, it is necessary to evaluate the relation of this condition to age and other relevant factors. From unpublished data collected by the Public Health Service, tabulations of lung-field markings in relation to age were made on two groups of men. One group comprised 229 men who had never been employed in a dusty trade; the other group included 80 men engaged in mining and refining kaolin by wet methods. The percentage of men who have a second-degree exaggeration of the linear pulmonic markings increases with advancing age. It is significant, however, that in the entire group of 309 men there were no cases of ground-glass lung-field markings. Other studies confirm this. Dreessen (43a) examined 80 men exposed to marble dust; 18 percent had second-degree linear markings, but none had ground-glass or nodular markings. Similar data are available (42) for 201 men employed in portland cement mills who were examined by Gardner and Sampson. No cases of granular markings or of discrete, nodular fibrosis were found.

In every age group the proportion of asbestos-exposed men who had advanced stages of pulmonary fibrosis is greater than in the men not exposed to asbestos dust. The percentage of men with ground-glass lung-field markings increases with advancing age. In view of the data for the two occupational groups just discussed and of the relation between this condition and length of employment, it appears that the reason why the incidence of this type of pulmonic markings is correlated with age is that the older people have been exposed to asbestos

TABLE 27.—*Relation of lung-field markings to age: Men who had never been employed in a dusty trade for the first time, kaolin workers, and men employed in asbestos textile factories*

Age	No previous dust exposure			Kaolin workers			Asbestos workers				
	Number of men	Percent with first-degree linear markings	Percent with second-degree linear markings	Number of men	Percent with first-degree linear markings	Percent with second-degree linear markings	Number of men	Percent with second-degree ground-glass markings	Percent with second-degree ground-glass and second-degree linear markings	Percent with first-degree ground-glass and second-degree linear markings	Percent with first-degree ground-glass and first-degree linear markings
Total examined	229	92	8	80	79	21	359	11	56	11	6
Under 20	58	93	7	7	100	0	11	73	27	0	0
20 to 29	107	91	6	29	93	7	109	20	16	7	4
30 to 39	38	90	10	23	65	35	112	22	24	17	6
40 to 49	17	88	12	12	60	40	76	40	25	21	13
50 to 59	7	57	43	6	50	50	47	55	28	12	21
60 to 69	2	100	0	3	100	0	2	50	0	0	50

FIGURE 26.—INCIDENCE OF GROUND-GLASS LUNG-FIELD MARKINGS ACCORDING TO LENGTH OF EMPLOYMENT AND DUST CONCENTRATION

TABLE 28.—*Proportion of asbestos textile workers with ground-glass lung-field markings*

Dust exposure (million particles per cubic foot)	Under 10	10 to 19	20 and over
Under 10			
Adjusted			
Expected			
Percentage			
10 to 19			
Adjusted			
Expected			
Percentage			
20 and over			
Adjusted			
Expected			
Percentage			

A21542


FIGURE 25.—RELATION OF GROUND-GLASS LUNG-FIELD MARKINGS TO DUST EXPOSURE

UCC 017707

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 34 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/34
UCC 017708
A21543

[illegible] Although it is probable that these [illegible] errors due to small numbers of exposed [illegible] fibers excite a different, and possibly [illegible] asbestos particles. If this were true, the [illegible] isolated above would not be a completely [illegible] measure of the pneumoconiosis-producing power of [illegible]

[illegible] unreasonable to suppose that the *amount* of pulmonary [illegible] more closely related to the weight of inhaled asbestos [illegible] other physical property of the dust. The number [illegible] used as a rough estimate of the weight of most [illegible] studies have shown that the two quantities are [illegible] This may not be as good a measure in the case



[illegible] 37.—RELATION OF DIAPHRAGMATIC FIXATION TO DUST EXPOSURE. THE [illegible] PERCENTAGE OF PERSONS IN EACH EXPOSURE GROUP WHO HAD FIRST, SECOND, [illegible] DEGREE DIAPHRAGMATIC FIXATION HAS BEEN REPRESENTED BY THE [illegible] OF VERTICAL BARS.

of asbestos because, as table 6 shows, there are many long fibers in the dust suspended in workroom air. The difficulties encountered in [illegible] long fibers have already been discussed (p. 23) and only one additional point need be mentioned here. A fiber 50 microns in [illegible] capable of reaching the lungs (table 37); it would be counted [illegible] particle in making dust counts, and yet it would have approximately 50 times the weight of a fragment of the same fiber only 1 micron in length. The 2 particles would have the same influence on the dust count, but one would carry 50 times as much asbestos to the lung as the other if both were inhaled and retained. Unfortunately, measurements of the weight of dust in the air were not made. If they had been, it is possible that the relation between asbestosis and dust exposure might have been altered.

During fluoroscopic examination of the chest the degree of movement of the diaphragm was observed and recorded. The percentage of people with impaired diaphragmatic movement increased with increasing length of dust exposure, and the proportion of workers with severe degrees of impairment likewise increased.

TABLE 29.—*Occurrence of diaphragmatic fixation in relation to million particle years of exposure*[1]

PERCENT

Degree of fixation	Controls	Million particle years			
		Under 25	25 to [illegible]	[illegible]	[illegible]
Normal	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
+	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
++	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
+++	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

NUMBER OF CASES

Degree of fixation	Controls	Million particle years			
		Under 25	25 to [illegible]	[illegible]	[illegible]
Total tested	76	195	[illegible]	55	[illegible]
Normal	74	172	[illegible]	32	[illegible]
+	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
++	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
+++	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

[1] The percentage values in the upper half of this table were obtained by dividing the values tabulated "total tested" by the corresponding values in the lower half of the table.

This is shown in figure 37 where asbestos workers have been divided into five groups (one of them a control group), according to the total amounts of asbestos dust to which they have been exposed during their working life. Only two of the 76 controls had any impairment of diaphragmatic movement. For each group the percentage with normal diaphragmatic movement, and slight (+), moderate (++), or severe (+++) diaphragmatic fixation has been calculated. These percentages have been represented by the heights of vertical bars.

Severe forms of diaphragmatic fixation accompanied advanced stages of lung fibrosis. Among 295 persons with first-degree linear lung-field markings, 9.5 percent had impaired diaphragmatic movement. Of the 22 persons with second-degree, ground-glass lung-field markings, 68 percent had impaired diaphragmatic excursion. This is a reasonable finding, since lung fibrosis and diaphragmatic limitation appear to be part of the same process.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 35 of 57
UCC 017709
A 21544
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/35

[illegible] pleural thickening, did not occur to any [illegible] workers although they are common among [illegible] study of granite workers carried on by the [illegible] "Irregularity of the diaphragm were [illegible] Sixty percent of cases had evidence of it, and as the [illegible] there was an increase in changes of the diaphragm [illegible] These were instances in which angulation of the [illegible] was in evidence and others which showed indications of [illegible]." In a study of anthracite miners carried on by [illegible] Health Service 51, p. 58): "Decrease in the motibility of [illegible] diaphragm as demonstrated by fluoroscopy was frequently observed. This malfunction, varying from slight to practical fixation, increased with the severity of the pulmonary pathology. Not infrequently adhesions between the visceral and diaphragmatic pleura could be seen." On the contrary, in asbestosis, with the exception of [illegible] (fig. 18), there was no roentgenographic evidence of diaphragmatic pleurisy. It appears that the limitation of movement, which is a definite part of the symptom complex of asbestosis, may be due to inelasticity of the lungs and binding of the lungs to the parietes and the mediastinal structures by fibrous pleuritis. (See pathological [illegible].)

PHYSICAL FINDINGS

The objective symptoms that are most frequently observed and most useful in establishing a diagnosis of asbestosis are general appearance of ill health (particularly any signs of loss of weight), the condition of the finger nails, and diminished chest expansion. These symptoms obviously appear late in the course of the disease. The useful chest signs seem to be limited to adventitious sounds. Diminished breath sounds, prolonged expiration and crackling râles over the bases of the lungs and intercapular regions seem to accompany asbestosis. Various râles and adventitious sounds were heard in the chests of 5 percent of the persons in the control group. Râles were noted in 14 percent of the asbestos-exposed persons who had normal lung-field markings or who had only first-degree linear exaggeration. However, in the group of persons with ground-glass lung-field markings (persons who had asbestosis), 49 percent had râles or other adventitious sounds. In about two-fifths of the cases, râles persisted after [illegible].

Changes in the percussion note are hard to evaluate, probably because the condition tending to give a dull note is counterbalanced by [illegible] emphysema which produces hyper-resonance. The presence of localized areas of conglomerate fibrosis and localized areas in which there is marked compensatory emphysema obviously accounts for the lack of more definite findings.

Chest expansion decreases with increasing dust exposure and with increasing pulmonary fibrosis. As the exposed men were divided into four groups according to dust exposure, and the average chest expansion and the average chest girth at expiration were calculated for each group. The same men were classified into four groups on the

PHYSICAL EXAMINATION

General Appearance..... Hair..... Eyes..... Ht..... Wt..... Gr. Wt.....

Skin and mucous membranes..... Nose..... Mouth and teeth.....

Eyes: Pupils regular..... Irregular..... React to light & A..... Glasses.....

Ears: Hearing..... Discharge..... Throat..... Tonsils..... Pharynx.....

Teeth..... Gums..... Pyorrhea.....

Glands: Thyroid, Normal..... Cervical..... Epitrochlear..... Axillary.....

Blood pressure..... Chest: Type.....

Exercise	Rate at rest	Exercise	After 1 minute rest
Pulse: regular			
irregular			
Respiration			


Expansion.....

Palpation.....

Circumf.: Inspiration.....

Expiration.....

Expansion.....



Heart: Size, Normal..... Enlarged..... Liver, Normal.....

Murmurs..... Aromatic odor.....

Hernia..... Enlarged Rings.....

Extremities: Clubbing..... Cyanosis..... Bowed nails..... Edema.....

Special marks of identification.....

Remarks.....

Examined by.....

FIGURE 38.—FORM USED FOR RECORDING PHYSICAL FINDINGS

basis of lung-field markings, and similar calculations were made. The decrease in average chest expansion cannot be attributed to any lack of large-chested men in the groups with high dust exposure or advanced pulmonary fibrosis. On the contrary, the average chest girth increases slightly with increasing dust exposure and with increasing pulmonary fibrosis. Several factors may have contributed to the

may appear to be hyperexplicit. In other words, the general appearance of the chest is that of the [illegible]

Among the conditions that may [illegible] are pectoral adhesions and the relative inelasticity of fibrous lung tissue.

Respiratory rate

The direct effect of pulmonary fibrosis on respiratory rate may be seen in table 33. Before the exercise test the respiratory rate of males with ground-glass lung-field markings was higher, but not significantly so, than the respiratory rate of males with first or second linear expansion of pulmonic markings. The exercise test consisted of having the subject stand with one foot on a box 11 inches high and to raise the other foot to the level of the top of the box 25 times in 30 seconds. The respiratory movements were counted during a 15-second period preceding the test, during a 15-second period immediately after exercise, and during a 15-second period 2 minutes after the test. Exercise raised the respiratory rate of all four groups of men, but the rise in respiratory rate was dependent upon the degree of pulmonary fibrosis. The men with ground-glass lung-field markings being affected least. After 2 minutes' rest, the fall in respiratory rate was again dependent on pulmonary fibrosis. The respiratory rate returned most closely to the prexercise rate in men with normal pulmonic markings or with first-degree linear exaggeration of lung-field markings. The return to the prexercise rate was slower in men with asbestosis or ground-glass lung-field markings).

TABLE 33.—*Relation of respiratory rate to lung-field markings in male asbestos workers*

Average respiratory rate	Lung-field markings			
	First-degree linear	Second-degree linear	Ground-glass lung-field plus	Second-degree ground-glass plus
Number of men examined	25	25	25	25
Before exercise	18.12	18.25	18.40	18.48
After exercise	23.72	24.80	24.80	24.88
2 minutes after exercise	20.12	20.16	20.20	20.28

TABLE 34.—*Influence of asbestosis on the respiratory rate of male workers*[1]

Average	Exposed to asbestos dust, roentgenologically negative	Asbestosis
Number of men examined	25	25
Rate before exercise	18.20	18.40
Rate after exercise	23.72	24.80
Rate after rest	20.12	20.16

[1] These data were not included in Public Health Bulletin No. 293.

Chest expansion

Chest expansion tends to decrease somewhat with advancing age, but the average chest expansion is less in asbestos-exposed men than in men of comparable age who have not been so exposed.

TABLE 35.—*Average chest expansion of males*

Number of men in age group	Exposed to kaolin		Employed in asbestos textile factories but not exposed to dust		Exposed to asbestos dust	
	Average chest expansion	Number of men in age group	Average chest expansion	Number of men in age group	Average chest expansion	
	Inches		*Inches*		*Inches*	
	3.69	4	4.25	9	3.25	
	3.92	35	3.54	15	3.55	
	3.45	19	3.11	10	3.05	
	3.29	5	2.80	8	2.95	
	3.17	3	2.81	6	2.95	
	2.88	3	2.60	2	2.80	

Essentially, the chest shows evidence of limited respiratory excursion; the supra- and infraclavicular fossae are conspicuous; the anteroposterior diameter of the chest is increased; the substernal angle may be more obtuse; evidence of wasting or loss of subcutaneous tissue may be present; certain of the accessory muscles of respiration

A21545

UCC 017710

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 37 of 57
UCC 017711
A21548
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/37

[illegible] anthracite miners. In [illegible] the incidence rate was definitely higher [illegible] normal men before exercise, after exercise, and [illegible].

[illegible] FINGERS AND CURVING OF NAILS

Many asbestos-exposed workers, particularly those with long dust [illegible], had clubbing of the fingers. The terminal phalanges may become so large that the finger resembles the end of a drumstick, a condition conveniently described by the term clubbed fingers. Clubbing is accompanied by a downward curving of the end of the nail.

The clubbing is almost always bilaterally symmetrical and often involves the toes as well as the fingers. It is common in pulmonary tuberculosis, in bronchiectasis, empyema, and congenital heart disease it may be seen in its most exaggerated form. Such exaggeration, however, was comparatively rare in this study. In a study of 2,711 [illegible] anthracite workers, the deformity, regardless of degree, was observed in about a third of the workers having anthraco-silicosis, but was present in only 1 percent of the controls. The real significance of the affection has not been determined. Norris and Landis (62) state that in all probability it is a manifestation of long-standing toxemia associated with phthisis or bronchiectasis. The bronchiectatic and emphysematous changes noted in the asbestotic lung at autopsy probably account to a great extent for the occurrence of the change in the workers with asbestosis.

TABLE 15.—*Number and percent of asbestos-exposed workers who had clubbed fingers or curved finger nails, classified by lung-field marking*

[illegible]	Total examined		Lung-field markings								Total
			No abnormal finding		Second-degree fibrosis		First-degree ground glass		Second-degree ground glass		
	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
[illegible]	[illegible]	[illegible]	78	100	50	100	22	100	163		
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

Five controls (6 percent) had curving of the nails. This percentage was much higher (19.2) in the asbestos-exposed group. Thirty-nine percent of the workers with first- or second-degree ground-glass lung-field markings had this condition.

HEART DEFECTS

Taking age into consideration, blood pressure values conformed closely to the values found on examination of 10,143 male industrial [illegible] (36).

Heart defects were no more numerous than [illegible] general. They were tabulated in relation [illegible] and pulmonary fibrosis in the same way as the [illegible] past dis-




FIGURE 39.—PHOTOMICROGRAPHS OF ASBESTOSIS BODIES FOUND IN SECTIONS ENLARGED 530 TIMES EXCEPT E WHICH IS ENLARGED 310 TIMES. A SERIES, ROLLED IN UNITS OF 50 MICRONS, HAS BEEN DRAWN BESIDE EACH [illegible] BODY.

cussed, but no simple relationships could be found. A careful cardiac study (including electrocardiograms) was made on one group of asbestos workers. Many workers with advanced cases of asbestosis had normal hearts and no evidence was found which would relate heart impairments to asbestos dust exposure.

may appear to be hypertrophied. In other words, the general appearance of the chest is that of the "all pie" type of physique. Among the conditions that may reduce chest expansion are pleural adhesions and the relative inelasticity of fibrous lung tissue.

MEASUREMENT OF CHEST EXPANSION

The direct effect of pulmonary fibrosis on respiratory rate may be seen in table 33. Before the exercise test the respiratory rate of males with ground-glass lung-field markings was higher, but not significantly so, than the respiratory rate of males with first- to third-degree linear evidence of pulmonic markings. The exercise test consisted of having the subject stand with one foot on a box 11 inches high and to raise the other foot to the level of the top of the box 25 times in 30 seconds. The respiratory movements were counted during a 15-second period immediately preceding the test, during a 15-second period immediately after exercise, and during a 15-second period 2 minutes after the test. Exercise raised the respiratory rate of all four groups of men, but the rise in respiratory rate was dependent upon the degree of pulmonary fibrosis, the men with ground-glass lung-field markings being affected most. After 2 minutes' rest, the fall in respiratory rate was again dependent on pulmonary fibrosis. The respiratory rate returned most closely to the preexercise rate in men with normal pulmonic markings or with first-degree linear exaggeration of lung-field markings. The return to the preexercise rate was slower in men with asbestosis (ground-glass lung-field markings).

TABLE 33.—*Relation of respiratory rate to lung-field markings in male asbestos workers*

Average respiratory rate	Lung-field markings			
	First-degree linear	Second-degree linear	Fine to coarse (ground-glass)	
Number of men examined				
Before exercise				
After exercise				
2 minutes after exercise				

TABLE 34.—*Influence of asbestosis on the respiratory rate of asbestos workers* [1]

Average	Exposed to asbestos dust, some fully treated	Exposed to asbestos dust, fully treated	Not exposed to asbestos
Number of men examined			
Rate before exercise			
Rate after exercise			
Rate after rest			

[1] These data were not included in Public Health Bulletin No. 231.

A21547

Data on the average chest expansion for males for [illegible] Since the trend is slight it may be [illegible] expansion to the girth increases with [illegible] with increasing pulmonary fibrosis.

Data on the average chest expansion in male asbestos workers

TABLE 35.—*Relation of average chest expansion to lung-field markings in male asbestos workers*

Dual expansion, millimeters per year	Lung-field markings			
	First-degree linear	Second-degree linear	First-degree ground glass	Second-degree ground glass
Less than 25				
25 to 49				
50 to 75				
More than 75				

Chest expansion tends to decrease somewhat with advancing age, but the average chest expansion is less in asbestos-exposed men than in a group of comparable age who have not been so exposed.

TABLE 36.—*Average chest expansion of males*

Number of men in age group	Average chest expansion	Employed in asbestos textile factories but not exposed to dust	Number of men in age group	Average chest expansion	Exposed to asbestos dust
	Inches			*Inches*	

Externally, the thorax shows evidence of limited respiratory movement; the supra- and infra-clavicular fossae are conspicuous; the anteroposterior diameter of the chest is increased; the substernal angle may be more obtuse; evidence of wasting or loss of subcutaneous fat may be present; certain of the accessory muscles of respiration

UCC 017712

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/38

A21548

FIGURE 105.—APPEARANCE OF THUMB OF 36-YEAR-OLD MAN EMPLOYED 4 YEARS IN CARD ROOM OF AN ASBESTOS TEXTILE FACTORY. PREVIOUSLY EMPLOYED 11 YEARS IN COTTON AND WOOLEN TEXTILE MILLS. PHOTOGRAPH [illegible] 2X.

UCC 017713

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 40 of 57
UCC 017714
A21549
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/40

[illegible] ; their [illegible] complete [illegible] series, [illegible] found since [illegible] to be a [illegible] indicate the [illegible] asbestosis bodies most [illegible] The largest body found in subsequent study of [illegible] measured 181 microns in length. There seems to [illegible] no satisfactory explanation for the absence of asbestosis [illegible] measuring less than 10 microns in length.

[illegible] appears to differ from other pneumoconiosis-producing [illegible] with respect to the size of the largest particle that may reach [illegible]. The largest reported upper limit of size for rock dust in [illegible] gold mines of South Africa and in the granite (48) and anthracite [illegible] asbestosis in this country is 14 microns. The data of table 37 are in harmony with a concept that has been gaining support in recent years, namely, that particles beyond 10 microns in size reach the alveoli. Measurements indicate that the diameter of air sacs may be from 150 to 300 microns in adults, or, in other words, of such a size that asbestos fiber or resulting asbestosis bodies may be readily accommodated.

Asbestosis bodies were found more frequently in the sputum of persons who complained of a persistent cough than in the sputum of persons without a cough. This is a reasonable finding, since the act of coughing tends to raise materials from the lungs. Half of the 50 people who complained of dyspnoea had asbestosis bodies in their sputum. Less than a fifth of the 225 asbestos-exposed people who did not make a complaint of dyspnoea had asbestosis bodies in their sputum. Almost the same relationship holds for a complaint of [illegible].

Asbestosis bodies were found in the sputum of 46.9 percent of persons whose condition was diagnosed as asbestosis. A much smaller proportion, 21.3 percent, of essentially normal asbestos-exposed persons had asbestosis bodies in their sputum. Thus, it appears that the presence of asbestosis bodies is evidence of exposure to asbestos dust, but it is not necessarily a diagnostic sign of asbestosis. Asbestosis bodies were found in the sputum of persons who had been exposed to asbestos dust for as short a time as 3 months. Of the workers employed less than 1 year, 22.5 percent had asbestosis bodies in their sputum. A larger percentage, 38 percent, of workers employed more than 5 years had these bodies in their sputum. In evaluating these percentages it must be remembered that they are based on the examination of a single sputum specimen from each individual. If several specimens could have been collected and examined, the percentage of persons with asbestosis bodies would almost certainly be much higher.

No asbestosis bodies were found in sputum samples from any of the [illegible] persons tested who were not exposed to asbestos dust.

The incidence of asbestosis bodies increased with [illegible] exposure.

TABLE 38.—*Relation of dust concentration to occurrence of asbestosis bodies*

Weighted dust exposure	Number exposed	Number with asbestosis bodies found	Percent with asbestosis bodies found
Total exposed	250	74	29.6
0 to 4.9	44	4	9.1
5 to 9.9	90	21	23.3
10 to 19.9	69	24	34.8
20 and over	47	25	53.2

RESULTS OF TESTS ON URINE

Each person employed in the three asbestos plants studied was asked to bring a 24-hour sample of urine in a metal container provided for that purpose. In addition to the standard clinical tests, the silica content was determined by the method of King and Dolan (39).[1] Of the 416 persons exposed to asbestos dust at the time of the study, urine analyses are available for 403, and these results have been kept separate from the results of urine tests made on 67 persons whose work did not expose them to asbestos dust.

Albumin was found in the urine of 10 of the 248 white males tested; in the urine of 10 of the 98 white females; and in the urine of 3 of the 57 negro males tested. Only one sample of urine (from a white male) contained sugar (Benedict's test). These rates of incidence are no higher than in the controls and, insofar as published data are available for comparison, they seem to be no higher than the rates generally found in industrial populations.

TABLE 39.—*Relation of urinary silica to dust exposure*

Urinary silica	Controls	Dust exposure, million particles per cubic foot			
		0 to 4.9	5 to 9.9	10 to 19.9	Over 20
Milligrams silica per liter	3.0±2.0 to 30.52±1.11	3.55±0.13 to 56.32±1.28	2.95±0.05 to 50.29±1.25	3.2±0.13 to 25.47±1.1	4.11±0.2 to 45.25±1.1
Milligrams silica per day					
Number tested	65	98	105	77	23

Silica was present in concentrations ranging from 0.7 to 11.1 mg per 100 cc. Average urinary silica concentrations for groups of persons whose dust exposure varied widely (table 39) do not indicate that there was any simple relation between the exposure to asbestos dust (which consists of hydrated magnesium silicate) and the concentration of urinary silica. Rather, it appears that the values for

[1] The colorimetric procedure for the determination of silica in the urine consists in the production of a silico molybdic acid complex which is reduced to give a blue color, the depth of color being proportional to the amount of silica present.

A21550


Experimental findings.—In a study in which [illegible] it is important to exclude the possibility that an observed change may have resulted from inhalation of another kind of dust or some other disease process. An early case of silicosis, for instance, might present signs and symptoms which would differ from those of asbestosis mainly in degree. A diagnosis of asbestosis should be made only when the occupational history shows previous exposure to asbestos dust and no heavy exposure to other siliceous dust.

X-ray findings.—Although the ground-glass markings that are so characteristic of asbestosis are usually localized in the basal portions of the lung fields, nevertheless all regions of the chest must be examined roentgenographically in order to detect signs of other disease processes that might complicate diagnosis. It is important, of course, to look for the signs of pulmonary tuberculosis. The distribution and the character of lung-field markings are significant. Shagginess of the heart shadow (see p. 50) has been mentioned by several physicians who have reported cases of asbestosis. The physical findings should be referred to whenever cardiac enlargement is found during the X-ray examination. A consideration of the chest shape of the individual will sometimes modify the interpretation of certain of these findings. Fluoroscopic examination is, of course, the best means of detecting diaphragmatic impairments, but valuable information can often be gained from a study of a chest X-ray film. Heavy pectoral muscles may cast an X-ray shadow which resembles in a single plate the shadow cast by the increased fibrosis in the bases. This possibility must always be kept in mind during the examination of chest X-ray films. Notes on the thickness of the chest muscles should be entered on the physical examination form, or these films should be made in order to obviate this source of error. The presence of bronchiectasis caused difficulty in diagnosing one case.

Classification of lung-field markings.—In this study, after examining an X-ray film and before making a diagnosis, the film was classified according to the nature and degree of fibrosis into one of four classes: first- or second-degree linear exaggeration of the lung-field markings or first- or second-degree ground-glass lung-field markings. There is good reason to believe that these four classes represent successive stages through which a typical case of asbestosis would pass under continuous exposure to asbestos dust. Since, as has been said, the ground-glass markings appear first in the basal portions of the lung fields, it follows that any given chest film might have areas in which the markings were ground glass and other areas in which the markings were linear. Usually the basis of classification has been the type of marking that predominates in the lower third of the visible lung field.

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/41

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 42 of 57



UCC 017716
A21551

FIGURE 40.—PORTIONS OF CHEST X-RAY FILMS WHICH ILLUSTRATE THE SUCCESSIVE CHANGES BROUGHT ABOUT BY LONG-CONTINUED INHALATION OF

In order to show these markings on a latter scale than would be possible if the entire film were reproduced and to illustrate this system of classification, figure 40 has been prepared.

Symptoms.—Dyspnoea, cough, blood-tinged sputum, pain in the chest, weakness, decreased chest expansion, curving of the finger nails and adventitious chest sounds are symptoms which also occur in silicosis. It is possible to differentiate pulmonary asbestosis from silicosis by history of exposure to asbestos, the presence of asbestosis[1] bodies in the sputum or lungs, and by the type and distribution of the X-ray markings. With reference to asbestosis bodies in the sputum, the data show that they may appear after 3 months' exposure. They show evidence of exposure to dust, but not established disease. Lynch and Smith (38) reported difficulty in demonstrating the bodies in an advanced case. The presence of rosettes as illustrated in figure 39 has been reported to be of greater diagnostic significance.

No advanced cases of asbestosis of the bed-ridden type were observed in the present study. Reports (13, 20, 26, 32) of fatal American asbestosis cases which were relatively uncomplicated by the presence of other disease processes are essentially similar to reports of uncomplicated British cases (9, 14, 34, 35). Patients in the terminal stage are usually reported to be emaciated, markedly dyspnoeic, cyanotic, and usually have a severe cough with wiry mucoid sputum.

REPRESENTATIVE CASE HISTORIES

Plates made from representative X-ray films appear as figures 41 to 51. Roentgenograms of a normal chest (figure 41) and of a chest showing advanced silicosis (figure 42) have been included for purposes of comparison. In figures 43 to 51 which represent asbestotic patients, an abstract from the case history and physical examination accompanies the X-ray. In viewing these roentgenograms it should be remembered that the ground-glass lung-field markings that are characteristic of asbestosis are exceedingly difficult to photograph and reproduce. It will be noted that the distribution of the shadows is not quite as general as in silicosis.

[1] Cooke has reported somewhat similar bodies in the sputum of anthracite miners, but they can be differentiated morphologically from asbestosis bodies.

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/42

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 43 of 57
UCC017717
A21553
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/43

...of table 3 provide a simple means of comparing asbestosis ... In a previously reported study of the ... it was found that workmen in anthracite ... could be divided into three groups which differed with respect ... the ratings of the dust to which they were exposed. The percent-

TABLE 43.—*Percentage of workers exposed to certain concentrations of asbestos dust in which extensive dust control measures are used to a limited extent and in which no effective dust control is practiced*

Dust concentration (particles per cubic foot)	Limited use of dust-control measures[1]	Extensive use of dust-control measures[2]
Over 10	18	7
5 to 10	28	15
2.5 to 5.0	35	32
Under 2.5	19	41

[1] Data from table 17.
[2] Data from (5).

ages of men having any stage of anthraco-silicosis (1, 2, or 3) have been tabulated separately for each group in relation to average dust concentration and length of employment. In passing, it may be of interest to call attention to the difference between the pneumoconiosis-producing power of the three kinds of dust encountered in anthracite mines. If one compares groups of men with the same dust exposure and the same length of employment, in general the percentage of men with anthraco-silicosis increases with the free silica content.

TABLE 45.—*Percentage of men having anthraco-silicosis*[1]

Dust concentration in million particles per cubic foot	1,115 anthracite miners exposed to dust containing less than 5 percent free silica			502 men employed in headageways who were exposed to dust containing about 13 percent free silica			151 rock workers exposed to dust containing about 35 percent free silica		
	Years in industry			Years in industry			Years in industry		
	Less than 15	15 to 24	Over 25	Less than 15	15 to 24	Over 25	Less than 15	15 to 24	Over 25
5.0 to 10.0	1.1	1.5	7.1	0.1	3.1	23.3	(2)	(2)	(2)
2.5 to 5.0	1.5	11.1	51.0	0	0	28.3	12.5	(2)	(2)
0 to 2.5	0	21.1	71.1	(2)	(2)	(2)	21.5	71.4	(2)
Over 10.0	4.7	58.1	88.8	(2)	(2)	(2)	5.8	56.2	62.0

[1] Cecil, Oliver, p. 50, Public Health Bulletin No. 221.
[2] Less than 10 men examined.

As in asbestosis, the incidence of pneumoconiosis among anthracite miners increases with increasing length of employment. There is a much more definite increase in the incidence of anthraco-silicosis with increasing dust concentration than there is in asbestosis, probably because men were exposed to a much greater range of dust concentrations in anthracite mines than in asbestos textile factories.

In comparison with the three kinds of dust encountered in anthracite mines, relatively low concentrations of asbestos dust and relatively short lengths of employment in the asbestos textile industry result in pneumoconiosis.

There was no free silica (quartz) in the dust encountered in the asbestos textile factories.

According to table 43, 16.3 percent of 69 former asbestos workers and 378 men and women employed at the time of the study in three asbestos textile factories had asbestosis. (Persons who were not exposed to appreciable quantities of asbestos dust have not been included.) Eight percent of the 378 workers employed at the time of study had asbestosis. This does not completely represent the incidence of asbestosis because about 150 employees were discharged from these factories 15 months before the present study was made. Many of the discharged asbestos workers had asbestosis, as the reports of Shull (31), Donnelly (12), and McPheeters (23) show. Although an attempt was made to examine as many as possible of these discharged cases, only 69 could be examined, 43 persons of whom had asbestosis according to the diagnostic standards employed in this study. If one assumes that the number of persons employed in these factories was the same before these 150 persons were discharged as it was when the study was made, and if one assumes that 62 percent of all the discharged cases had asbestosis, then 50 cases of asbestosis escaped observation. Adding these to the 73 that were found, dividing the sum by 447, and multiplying by 100 gives an estimated percentage of 27.5. This is not an unusual percentage value. In numerous studies of the dusty trades the percentage of workers with pneumoconiosis has been found to be between 20 and 25 percent. Two such studies may be mentioned. Of 7,722 miners and men seeking employment in lead and zinc mines at Picher, Okla., who were examined in 1927–28 (49), 21.3 percent had silicosis. In a study of anthracite miners (51), "The prevalence of anthraco-silicosis among the entire group of employees was found to be about 23 percent."

Other investigators have reported the incidence of asbestosis in asbestos textile factories. Lanza, McConnell, and Fehnel (17) reported, "Of the total of 126 X-ray examinations, 4 were diagnosed as second-degree asbestosis, 63 as first degree, 39 as doubtful, and 20 as negative." The persons he studied "were selected more or less at random from among those having more than 3 years of employment in the industry." Since Lanza did not examine all the employees in the factories he studied, percentage values calculated from his data are not directly comparable with rate of incidence for the present study.

[illegible] Merewether, and Pontz (15) noted that "Asbestosis [illegible] 4 persons (2 men, 2 women), or 25 percent of the [illegible] Queen

INCIDENCE OF ASBESTOSIS IN DIFFERENT OCCUPATIONS

The percentage of persons affected by asbestosis varies in different occupations. The numbers of persons employed in many of the occupational groups are so small that caution must be exercised in making comparisons, but the data seem to be of sufficient interest to warrant publication. Classified by departments, the preparation department has the highest percentage of asbestotic persons. Carding is the principal occupation in this department. Although only 8 of the 72 carders had been employed more than 10 years, 29.2 percent had asbestosis. Eleven carders had advanced asbestosis. No other occupational group had so large a proportion of advanced cases.

TABLE 46. *Number and percentage of workers having asbestosis, classified according to department and occupation*

By principal occupation	Number employed	Percent with asbestosis	Number with asbestosis		
			Stage 1	Stage 2	Stages 3 and 4
[illegible]	102	25	[illegible]	[illegible]	[illegible]
[illegible]	5	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	5	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	30	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	72	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	256	11	[illegible]	[illegible]	[illegible]
[illegible]	85	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	64	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	73	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	4	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	29	14	[illegible]	[illegible]	[illegible]
[illegible]	13	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	11	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	15	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	18	11	[illegible]	[illegible]	[illegible]
[illegible]	15	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	3	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]	29	3	[illegible]	[illegible]	[illegible]
[illegible]	17	0	[illegible]	[illegible]	[illegible]
[illegible]	31	0	[illegible]	[illegible]	[illegible]
Total	502	10	25	25	21

In the spinning, twisting, and winding department, the mule and ring spinners have a higher proportion of asbestotic persons than the average, 24.6 percent. Twenty of the sixty-five mule and ring spinners had been employed more than 10 years.

A third occupation with an excessive prevalence rate for asbestosis is cloth weaving, in which 25.6 percent of the workers are affected. Only 3 of the 43 workers had been employed more than 10 years.

UCC 017718

A21560

ASBESTOSIS IN GREAT BRITAIN AND IN THE UNITED STATES

Lanza (18) has called attention to a point of interest, namely, that British investigators have found asbestosis more severe and menacing than it was in the United States. He adds that this difference is probably more apparent than real. Lynch and Smith (20) have reported 3 fatal cases whose duration of employment in asbestos textile mills was 11½, 17, and 21 years in occupations having dust exposures of the order of dry weaving. Stock (32) reported a fatal case in a man 27 years of age who worked 8 years as a carder in an asbestos plant. Egbert (13) has also given complete findings of similar case in a man 30 years of age who had worked 13 years as a carder and who died 4 years after onset of symptoms and 3½ years after discontinuing work.

These examples of American fatalities, some in young individuals, suggest that the period of dust exposure before disability or death ensues is as short as is the case in silicosis.

Bridges (63) in the factory inspector's reports of Great Britain has shown from year to year (table 47) that the duration of employment of fatal cases with asbestosis is only half that of cases with silicosis.

TABLE 47.—*Silicosis and asbestosis in Great Britain*

	Number of deaths	Average age at death	Duration of employment		
			Mean	Min.	Max.
Through 1915:					
Silicosis	311	[illegible]	[illegible]	[illegible]	[illegible]
Silicosis with tuberculosis	772	[illegible]	[illegible]	[illegible]	[illegible]
Asbestosis	52	[illegible]	[illegible]	[illegible]	[illegible]
Asbestosis with tuberculosis	46	[illegible]	[illegible]	[illegible]	[illegible]

More than 90 percent of the raw mineral used in the manufacture of asbestos textiles in this country and Great Britain is Canadian chrysotile, indicating that very little difference is to be expected from the standpoint of biological reaction in the two countries. As far as dust concentrations in the various operations of asbestos manufacture of British practice can be compared with results obtained in this country, no greater differences are found than exist in the inherent limitations of the dust-sampling devices. It would seem, therefore, that quite negligible differences exist in the manifestations of the disease here and abroad with the possible exception of associated or complicating tuberculosis.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 44 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/44

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 45 of 57

PATHOLOGICAL FINDINGS IN ASBESTOSIS

Specimens for pathological study were furnished through the courtesy of Dr. J. Rush Shell, Charlotte, N. C., and Dr. H. F. Easom, director of the Division of Industrial Hygiene of the North Carolina State Board of Health.

The pathology of asbestosis is characterized by diffuse, interstitial, pulmonary fibrosis and by the presence of asbestosis bodies in the lungs.

GROSS APPEARANCE

In gross appearance the lungs vary in color from brownish to pale gray. Usually, they are not distorted like silicotic or anthracosilicotic lungs unless an extensive amount of long-standing fibrosis is present. Often the lungs are firm and heavy, but softer areas can sometimes be felt on palpation. Crepitus depends on the amount of fibrosis. The pleura may be thickened and fibrous tags may be scattered over the surface. In the cases described in this report, thickening of the pleura and the presence of fibrous adhesions are more frequently found on the right lung than on the left. Adhesions to the diaphragm are sometimes present.

On section, small, dark gray, irregularly shaped areas are seen on the cut lung surface. They are rather evenly distributed and they are of approximately the same size. Visible strands of connective tissue of varying size run through the lung parenchyma, forming a more or less regular, fibrous network, and enclosing the dark gray, nodular areas and lung tissue which show alveoli. Areas of well-marked emphysema are noted in the apices and outer portions of the lower lobes, a feature that was referred to in the discussion of the X-ray findings. The bronchi and bronchioles appear to be distended and their walls are often thickened. In one case, this bronchiectasis is most marked in the upper lobe, particularly in the apex.

Congested areas are occasionally seen on gross inspection of some of the lung sections. They are probably terminal.

The hilar lymph nodes are not markedly increased in size. They are pinkish gray and contain about the usual amount of anthracotic pigment found in a city dweller.

MICROSCOPIC APPEARANCE

The pleura, when thickened, is made up of adult fibrous connective tissue cells. The collagen fibers are usually well developed and may show hyalinization. In some sections the thickened pleura is vascular,

UCC 017719

the blood vessels generally being small capillaries. Areas of variable size, containing anthracotic pigment, are occasionally found in the subpleural connective tissue.

Throughout the parenchyma of the lung a diffuse interstitial fibrosis is present, which varies with the asbestos-dust exposure of the individual. The areas of fibrous tissue are irregular or stellate in shape and may contain small blood vessels. The cellular elements consist of adult fibrous tissue cells, fibroblasts, some collagen, and often numerous macrophages. The strands of fibrous tissue represent an increase in the interlobular connective tissue. The fibrosis is usually rather uniformly distributed throughout the entire lung. In areas of considerable fibrosis, atelectasis is noted due to the contraction of the fibrous tissue. The alveoli occurring between areas of fibrosis are generally distended and show a moderate to marked emphysema. The alveolar walls are thickened by fibrous tissue, and in some sections hyalinization and the loss of nuclei occur. The alveoli are lined by a single layer of epithelial cells. The bronchial walls and the walls of the bronchioles may be thickened, and a slight to moderate lymphocytic infiltration is present occasionally. The smaller bronchi and bronchioles are frequently quite distended. The walls of the small blood vessels are thickened, particularly the adventitia, which contains an increased amount of fibrous tissue. Sections through emphysematous blebs show a fairly thick, fibrous, connective-tissue wall, lined by a single layer of flat or low cuboidal, epithelial cells. The blebs, measuring as much as 7 mm in diameter in the specimens studied, are apparently of bronchial origin.

A 21561

Asbestosis bodies are found in almost all parts of the pulmonary tissue. They occur most frequently in the areas of fibrosis and are accompanied by a greenish black pigment. They are also noted in the connective tissue about the small bronchi, in the alveolar walls, and in the alveoli. Macrophages and giant cells are usually present when the asbestosis bodies are found in the alveoli and often surround or engulf the bodies. These cells also contain a considerable amount of anthracotic pigment, and the dark greenish black pigment commonly found in asbestosis. A brownish yellow pigment in lesser amounts is also present. The asbestosis bodies occur in the lungs in a variety of shapes. The most frequent is a beaded rod with large rounded ends. Spindle shapes with beaded centers are also noted. Rosette formation is relatively frequent. There appears to be no relation between the number of asbestosis bodies and the amount of pulmonary fibrosis. In one case in which there is exceedingly little fibrosis, the asbestosis bodies are particularly numerous, while in another case, showing extensive fibrosis, they are quite infrequent. These bodies are golden yellow in color and are stained blue with potassium ferrocyanide, showing the presence of ferric iron. With this

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/45

stain some fine blue strands have been noted which are apparently asbestos fibers in the process of becoming asbestosis bodies. By the use of polarized light, fine fibers are often demonstrated in the tissue which are identified as asbestos. These may be seen in the peribronchial connective tissue, the interstitial connective tissue, the

FIGURE 53.—ASBESTOSIS BODY IN INTRA-ALVEOLAR CONNECTIVE TISSUE. PRACTICALLY NO FIBROSIS IN THIS AREA. CASE H-21. MAGNIFICATION 1,000X. (ARMY MEDICAL MUSEUM NO. 52255.)

alveolar walls, and in the alveolar spaces. The blue staining fibers and the asbestosis bodies are not visible when viewed with crossed Nicol prisms. Other fine crystalline material is sometimes noted.

The presence of complicating tuberculosis is often difficult to

A21562

FIGURE 52.—PHOTOMICROGRAPHS OF LUNG SECTIONS SHOWING THE PRESENCE OF NODULAR FIBROSIS AND EMPHYSEMA IN A SILICOTIC LUNG AND DIFFUSE FIBROSIS AND EMPHYSEMA IN AN ASBESTOTIC LUNG. A SECTION OF A NORMAL LUNG IS SHOWN FOR COMPARISON. MAGNIFICATION 16X.

UCC 017720

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 47 of 57

resemble the developing fibrous areas of asbestosis to a marked degree. It is not easy in many cases to differentiate caseation from old hyalinized connective tissue in the stained specimens. In one of the cases studied, the presence of acidfast bacilli in the alveoli and connective tissue of the alveolar walls without the presence of caseation was observed. This could possibly be an early tuberculous infection in the preinvasive stage, or it could well be a contaminating nonpathogenic acid-fast bacillus.



FIGURE 54.—ASBESTOSIS BODIES ACCOMPANIED BY GIANT CELLS IN AN AREA OF FIBROSED LUNG TISSUE. CASE H-21. MAGNIFICATION 705X. (ARMY MEDICAL MUSEUM NO. 52255.)

Samples of three varieties of asbestos (chrysotile, crocidolite, and

UCC 017721

using a technique developed by the Public Health Service for studying the physiological behavior of dust. This method has been described elsewhere (60). All three samples of asbestos dust produced an inert type of reaction in the peritoneal tissue. On the basis of this response, asbestos dust has been considered to be potentially harmful, especially when inhaled over long periods of time. Asbestos dust, like talc (43, 44), which also produces an inert reaction, causes a diffuse, pulmonary fibrosis as has been observed on clinical and post-mortem examination of persons exposed to these dusts.

SELECTED AUTOPSY CASE REPORTS

Case H-21

(Figs. 53 to 58)

Male, white, aged 28 years.

Cause of death.—Myocarditis.

Occupational history.—Worked for 12 years in the weaving room of an asbestos plant. Prior to this he had worked in a cotton mill.

Medical history.—Confined to bed for the most part of his last 6 months of life because of weakness, dyspnoea, and unproductive cough.

Pathological examination showed well-defined advanced asbestosis, evidenced by much diffuse interstitial pulmonary fibrosis, accompanied by asbestosis bodies, emphysema, bronchiectasis, and bronchiolectasis.

Chemical examination of lung tissue.—Sample from right lung—ash 3.3 percent, silica 0.6 percent, silica in ash 18.32 percent. Sample from left lung—ash 3.8 percent, silica 0.76 percent, silica in ash 20.13 percent.

A21563

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/47




FIGURE 50.--CASE H-21. LEFT LUNG. PLEURA MUCH THICKENED. FEWER FIBROUS ADHESIONS THAN WERE NOTED ON THE RIGHT LUNG.

43109—39——8

A21564




FIGURE 51.--CASE H-21. RIGHT LUNG. MARKEDLY THICKENED PLEURA WITH FUSION OF INTERLOBAR PLEURA. DENSE FIBROUS TAGS OVER ENTIRE SURFACE.

UCC 017722



FIGURE 58.- CASE H-21. SECTION OF LEFT LUNG. DISTENDED BRONCHI AND AIR SPACES AT APEX. RATHER UNIFORMLY DISTRIBUTED FIBROUS AND EMPHYSEMATOUS AREAS THROUGHOUT REST OF LUNG.

A21565



FIGURE 57.- CASE H-21. SECTION OF RIGHT LUNG. ADVANCED ASBESTOSIS. DISTENDED BRONCHI AND AIR SPACES AT BOTH APEX AND BASE. DIFFUSE FIBROSIS THROUGHOUT LUNG SUBSTANCE AND THICKENED PLEURA.

UCC 017723

FIGURE 42.—SILICOTIC PULMONIC MARKINGS. WHITE MALE, AGE 61.

Note coarse nodular appearance, with beginning coalescence of the shadows in the upper half of the left. Moderate emphysema both bases and slight nodular final distortion. Nodular markings of the type here shown were not noted in asbestos employees.

A21552

FIGURE 41.—NORMAL PULMONIC MARKINGS. WHITE MALE, AGE 23.

Note the linear type of pulmonic markings usually seen in the chest roentgenogram of an average industrial worker not exposed to a dust hazard. This man had worked 3 years in the decorating department of a pottery and 2 years at general outdoor labor.

UCC 017724




FIGURE 43.—EARLY ASBESTOSIS. WHITE MALE; AGE 29.

Occupational history.—Carder (asbestos), 4½ years; nondusty trades, 5½ years.
Estimated dust exposure.—(*a*) 35.3 million particles per cubic foot. (*b*) Million particle years, 159.
No significant past medical findings or complaints.
Physical examination.—Intermediate, slender white male. Height, 67¾ inches; weight, 142 pounds. Rather long chest; expansion, 2 inches. No change in fremitus, breath sounds, or percussion note. Blood pressure, 168/118. Pulse rate, 96, 120, and 80 before, immediately after, and 2 minutes after functional exercise. Urinary silica, 5.2 mg per 100 cc (24-hour output, 34.3 mg). Routine urinalysis negative; specific gravity 1.022.
X-ray.—First-degree ground-glass appearance not quite obliterating the coarse linear markings. Note the fine, close-set reticular markings just off each hilum and the apices remaining quite clear.
Diagnosis.—Early asbestosis without symptoms complicated by hypertensive heart disease.




FIGURE 44.—EARLY ASBESTOSIS. WHITE MALE, AGE 33.

Occupational history.—Mule spinner (asbestos), last 15 years; roving attendant (asbestos), 1 year; nondusty trades, 8½ years.
Estimated dust exposure.—(*a*) Weighted average, 10.5 million particles per cubic foot. (*b*) Million particle years, 168.2.
Past medical.—Shrapnel wound right parietal region of head, October 1918. Bilateral pneumonia as a child.
Complaints.—Slight weakness.
Physical examination.—Intermediate build, fairly well nourished. Height, 6[illegible] inches; weight, 144 pounds. Chest expansion, 3¼ inches. Blood pressure, 136/82. Pulse rate, 83, 120, and 84, and respiratory rate, 16, 24, and 18 (respectively before, immediately after, and 2 minutes after functional exercise test). Intermediate chest type with slight increase in A–P diameter of chest. Few nonpersistent crackling râles at right base. Asbestos corn on palmar surface of left wrist. Slight cyanosis of lips and nails. A dusky red cast to ears. Sputum negative for *B. tuberculosis* and asbestos bodies. Urinary silica, 3.6 mg per 100 cc (24 hours, 21.5 mg).
Fluoroscopy.—Moderate (3+) limitation of diaphragmatic excursion. Hila moderately increased in size and density.
X-ray.—Note the first-degree grainy or ground-glass appearance, not obliterating the linear markings, giving general reticular appearance to lung fields. Generalized fibrosis 1 +.
Diagnosis.—Asbestosis 2.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 51 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/51

FIGURE 46.—MODERATELY ADVANCED ASBESTOSIS. WHITE MALE, AGE 40.

Occupational history.—Mule spinner (asbestos), last 13 years; somebody trains, 4½ years; idle, 4½ years.
Estimated dust exposure.—(a) Weighted average, 13.5 million particles per cubic foot. (b) Million particle years, 175.7.
Past medical.—Influenza and pneumonia, 1918. Frequent disabling colds.
Complaints.—None.
Physical examination.—Intermediate white male. Height, 70½ inches; weight, 150 pounds. Intermediate chest shape; expansion, 2 inches. Dry, musical rales over bases of lungs. Urinary sliea, 5.6 mg per 100 cc (24-hour output, 80 mg).
Sputum not examined.
Fluoroscopy.—Showed slight limitation of diaphragmatic excursion, right and left with first-degree diffuse ground-glass appearance of the lung fields.
X-ray.—Shows first-degree granular appearance and obliterating the linear markings. It illustrates the fine reticular fibrosis associated with this condition and the tendency to be most dense at the bases of the lungs.
Diagnosis.—Asbestosis 3.


FIGURE 45.—MODERATELY ADVANCED ASBESTOSIS. WHITE MALE, AGE 46.

Occupational history.—Broadcloth weaver (asbestos), 7 years 5 months. Previously in nonasbestos trades, 17½ years including 10 years as farmer.
Estimated dust exposure.—(a) Weighted exposure, 18 million particles per cubic foot. (b) Million particle years, 126.2.
Past medical.—Tuberculosis, 1915. Pleurisy, right, March 1926.
Complaints.—Dry, hacking morning cough for 3 years productive thick mucoid sputum. Slight dyspnoea, 2 to 3 years. Blood-tinged sputum.
Physical examination.—Height, 70½ inches. Weight, 160 pounds. Blood pressure, 140/82. Pulse rate, 72. RS, and RS, equal; respiratory rate, 24, 28, and 24 (before, immediately after, and 2 minutes after functional exercise test). Patient complains of breathlessness after functional exercise. Chest expansion, 1 inch. Shoulders somewhat rounded. Crackling rales heard over both lungs on quiet breathing, and these persist after cough; 2+ curved finger nails; 3.1 mg silica per 100 cc (single specimen).
Fluoroscopy.—Left side of diaphragm moves more freely than right; 2+ limitation of excursion, especially on right.
X-ray.—Shows second-degree ground-glass appearance obliterating usual linear pulmonic markings. It also shows preponderance of fibrosis in the bases of lung and shaggy appearance of heart shadow.
Diagnosis.—Asbestosis 3 (moderately advanced).

A21554

UCC 017726

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 53 of 57
asbestosclaims.law/library/18707/53
Published by AsbestosClaims.Law



UCC 017727

FIGURE 47.—MODERATELY ADVANCED ASBESTOSIS. WHITE MALE, AGE 49.

Occupational history.— Mule spinner (asbestos), 25 years, working 10 to 11 hours a day; nondusty trades, 9 years.

Estimated dust exposure.—(*a*) Weighted average, 7.0 million particles per cubic foot. (*b*) Million particle years, 221.2.

Past medical.—Typhoid fever at age of 11. Influenza, 1918. Dry pleurisy, right in 1931; second attack in 1935 on left. More than average number of chest colds each year. Kidney colic, 1931.

Complaints.— Slight dyspnoea, especially in climbing stairs. Slight morning cough past 2 to 3 years productive of small amount of white frothy sputum.

Physical examination.—Tall, fairly well-developed man. Height 70 inches, and weight, 172 pounds. Blood pressure, 140/80. Pulse rate, 80, 100 and 80; respiratory rate, 21, 32, and 24 (respectively before, immediately after, and 2 minutes after functional exercise test). Chest is long, and well developed with some flaring of the costal margins. Chest expansion, 1½ inches. Numerous crackling rales at bases on deep breathing and after cough. Sputum (single specimen) was negative for acid-fast bacilli and asbestosis bodies. Urinary silica, 3.2 mg per 100 cc (24-hour output, 23.0 mg).

Fluoroscopy.—Diaphragm almost fixed and lung fields interpreted as showing a first-degree ground-glass appearance.

X-ray.—Note the fine grainy or ground-glass appearance fairly well limited to the lower two-thirds of each lung field.

Diagnosis.—Asbestosis 3.




FIGURE 48.—MODERATELY ADVANCED ASBESTOSIS. WHITE MALE, AGE 39.

Occupational history.—Broadcloth loom room (asbestos), 14 years; nondusty trades, 7 years.

Estimated dust exposure.—(*a*) Weighted average, 49.7 million particles per cubic foot. (*b*) Million particle years, 459.7.

Past medical.—Bilateral pneumonia as a child. Sinus disease for the past 2 years. Typhoid fever, 1911.

Complaints.—Dry, hacking, morning cough productive of 2 ounces of white, viscid sputum, past 2 years. Shortness of breath, 2½ years.

Physical examination.—Fairly well-developed, slender, white man. Height, 72 inches; weight, 163 pounds. Blood pressure, 104/70. Pulse rate, 68, 76, and 78. Respiratory rate, 21, 30, and 30 (before, immediately after, and 2 minutes after functional exercise test). Chest is long, slender type. Expansion, 1¾ inches. Heart sounds are distant with a pulmonic second sound somewhat accentuated.

Fluoroscopy.—Peaking of the diaphragm both sides. 3 cm excursion upon forced respiration.

X-ray.—Note the deformity of diaphragm. This case is the only one showing definite deformity in the group. The lung fields were interpreted as showing second-degree diffuse grainy or ground-glass appearance with emphysema.

Diagnosis.—Moderately advanced asbestosis (asbestosis 3).

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 54 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/54


UCC 017728
42156

FIGURE 49.—ADVANCED ASBESTOSIS. WHITE MALE, AGE 48.

Occupational history.—Male spinner (asbestos), 17½ years (1½ years in same room with carding machines; 11 years he worked 11 hours a day; nondusty trades, 17 years.

Estimated dust exposure.—(a) Weighted average, 10.6 million particles per cubic foot. (*b*) Million particle years, 103.8.

Past medical.—Frequent chest colds. Aerophagia for several years. Malaria about 1926. Influenza, 1929 and 1932.

Complaints.—Cough, 6 years' duration, possibly somewhat improved since quitting work in asbestos. Severe paroxysms of cough in morning, causing him to vomit frequently and productive of small amount of tenacious mucoid sputum. Moderate dyspnoea since 1928, unchanged. Lost 15 to 20 pounds in weight and associated weakness before quitting work in asbestos.

Physical examination.—Slender fairly well-developed and nourished white male. Height, 68 inches, weight, 131 pounds. Rather long narrow chest. Chest expansion, 1½ inches. Tactile and voice fremitus apparently normal. Slight impaired resonance lower half of each lung. Numerous persistent dry crackle mostly over lower portions of lungs. Faint breath sounds lower one-half right and left; 1+ curving of finger nails. Blood pressure, 135/80. Pulse and respiratory rate, respectively, 72, 100, and 68 and 28, 32, and 28, (before immediately after, and 2 minutes after the functional exercise test). Moderate dyspnoea following this exercise. Urinary silica (single specimen), 2 mg per 100 cc. Sputum negative for acidfast bacilli and asbestosis bodies.

Fluoroscopy.—2+ log of diaphragmatic excursion right and left. Rather poor radiolucency lower lung fields.

X-ray.—Shows second-degree grainy appearance with almost complete obliteration of linear markings. Generalized fibrosis 2+.



FIGURE 50.—ADVANCED ASBESTOSIS. WHITE MALE, AGE 55.

Occupational history.—Preparation department (asbestos), 11 years; nondusty trades, 27 years.

Estimated dust exposure.—(a) Weighted average, 38.0 million particles per cubic foot. (*b*) Million particle years, 545.

Past medical.—Maxillary sinuses drained in 1930. Typhoid fever as a child.

Complaints.—Dry cough, 10 years, productive of 2 ounces of white, thick sputum over a period of 24 hours. Rather severe dyspnoea, 3 years. He must walk very slowly to prevent dyspnoea. Occasional edema of ankles for the past 4 years.


Physical examination.—Height, 66 inches. Weight, 132 pounds. Blood pressure, 112/70. Moderate arteriosclerosis. Pulse rate, 72, 80, and 68. Respiratory rate, 48, 54, and 54 (before, immediately after, and 2 minutes after functional exercise test). Patient stopped on completion of about one-half of prescribed exercise, but expressed a willingness to continue. Chest expansion, 2 inches. Breathing is labored and shallow. Prolonged expiration. Fine crackles appear over the lower half of both lungs after coughing. Heart is enlarged to the left with the aortic second sound somewhat more prominent than the pulmonic second. First sound at the apex is split and is particularly snapping elsewhere; 2+ curving of the finger nails.

Fluoroscopy.—Aortic shadow is definitely widened. The cardiac shadow shows 1+ increase in size. Diaphragm on ordinary breathing moves a scant 1 cm; 3 cm on forced excursion.

X-ray.—Shows diffuse ground-glass appearance. Note that most of the distinctly linear markings have been obliterated.

Diagnosis.—Advanced asbestosis.

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 55 of 57



UCC 017729

FIGURE 31.—ASBESTOSIS WITH FIBROID TUBERCULOSIS. WHITE MALE. AGE 38.

Occupational history.—Mule spinner (asbestos), 11 years 8 months; nondusty trades, 12 years.

Estimated dust exposure.—(*a*) Weighted average, 7.9 million particles per cubic foot. (*b*) Million particle years, 88.5.

Past medical.—Received sanatorium treatment for tuberculosis 8 months in 1930. Had pneumonia as a child. Appendectomy in 1916.

Complaints.—Dry cough past 5 years, worse at night. Slight dyspnoea, 8 years.

Physical examination.—Very thin man. Weight, 131 pounds. Blood pressure, 100/70. Pulse rate, 72, 80, and 68. Respiratory rate, 16, 16, and 20 (before, immediately after, and 2 minutes after functional exercise test). Asthenic type of chest. Some deformity of chest due to scoliosis. Chest expansion, 1½ inches. Breath sounds are slightly increased over the extreme right upper. Very few dry rales right supra-scapula fossa. Heart: The pulmonic second sound is much greater than the aortic second.

Fluoroscopy.—Diaphragm moves freely and equally on both sides. On forced respiration, excursion is 3 cm on right and 4 cm on left.

Diagnosis.—Moderately advanced asbestosis with arrested fibrotic tuberculosis

INCIDENCE OF ASBESTOSIS

Preceding sections of this bulletin have stressed the importance of the X-ray findings in establishing a diagnosis of asbestosis. The reason, of course, is that X-ray examination is the most direct means now available for detecting the primary effect of inhaled asbestos dust on the body; namely, initiation of interstitial fibrosis of lung tissue. Classification of asbestosis into a series of stages which follow one another in their development is a difficult matter because it depends to a great degree on the interpretation of relatively minor changes in the lung-field markings. Other investigators have encountered the same difficulty. Lanza, McConnell, and Fehnel (17) and Fulton, Dooley, Matthews, and Houtz (15) set up a category of doubtful cases of asbestosis. Stage 1 in the accompanying tabulation more or less corresponds to their category of doubtful cases. It comprises cases in which diagnosis rests largely on X-ray evidence. Stage 2 represents early cases of asbestosis, and stage 3 designates moderately advanced cases. More advanced cases (stage 4) are relatively infrequent. Stage 5 was reserved for bed-ridden cases.

TABLE 41.—*Relation of asbestosis to lung-field markings*

X-ray findings	Asbestosis absent	Stages of asbestosis				
		1	2	3	4	5
Total	372	21	26	16	7	10
First-degree linear	297	.....	.....	.....	.....	5
Second-degree linear	65	11	3	.....	.....	1
First-degree ground glass	6	13	22	8	.....	1
Second-degree ground glass	4	.....	1	8	7	3

Table 41 illustrates the relationship of stages of asbestosis to the X-ray findings. Although diagnoses of early stages of asbestosis were occasionally made in the absence of definite ground-glass lung-field markings when cough, dyspnoea, and objective findings were present in sufficiently advanced degree, diagnoses of the later stages of asbestosis were not made unless unmistakable ground-glass markings were present in the lung fields.

The incidence of asbestosis increases rapidly with increasing dust exposure. A fifth of the workers with an exposure of 50 to 99 million particle years, and half of the workers having an exposure of more than 100 million particle years had asbestosis. In comparison with anthraco-silicosis these are exceedingly low dust-exposure values

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/55

Union Carbide trial exhibit UC-4095: A Study of Asbestosis in the Asbestos Textile Industry, by Dreessen, Dallavall...
ACL Doc 18707 · Page 56 of 57
Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/56

UCC 017730
A21558

[illegible] of asbestosis to dust concentration and length of employment, in table 43, is of great practical importance because it [illegible] working conditions are most likely to cause asbestosis [illegible] long a period of exposure may elapse before the disease [illegible].

TABLE 42.—*Relation of asbestosis to total dust exposure* [illegible]

Diagnosis	Million particle years				Total
	Less than 25	25 to 49	50 to 99	Over 100	
[illegible]	121 5 0 0 0	81 2 0 0 0	43 5 0 0 0	12 12 23 11 7	257 24 23 11 7
[illegible]	126	83	50	105	322

[illegible] except [illegible] estimated and 2 persons with active pulmonary tuberculosis have [illegible] from the table.

TABLE 43.—*Occurrence of asbestosis in relation to dust concentration and length of employment*

PERCENTAGE WITH ASBESTOSIS

[illegible] exposure, million particles per cubic foot		Years in asbestos industry		
		0 to 4.9	5 to 9.9	Over 10
[illegible]	Affected Exposed Percentage	2 81 2%	1 19 5%	0 3 0%
[illegible]	Affected Exposed Percentage	0 70 0%	6 35 17%	11 19 58%
[illegible]	Affected Exposed Percentage	0 134 0%	22 44 50%	21 30 70%

The percentage of persons with asbestosis increases greatly with increasing length of employment. Lanza, McConnell, and Fehnel (17) reported a similar trend; their percentage values are higher than these. There are two explanations for the small number of persons employed 15 years or more. This group of asbestos textile factories has been in operation from 6 to 16 years, and because of the relatively short life of the industry in this section of the country comparatively few persons have been exposed to asbestos dust for a long time. Another explanation is suggested by the fact that almost three-fourths of the workers employed more than 15 years have the early stages, at least, of a disabling disease, asbestosis. Since there is no satisfactory way of locating and examining persons who drop out of an industry because of disability incurred therein, the percentage values of the foregoing tables are necessarily minimal estimates of the prevalence of

The incidence of asbestosis increases with increasing dust concentration. In their report of a similar study carried on in Pennsylvania asbestos textile factories, Fulton, Doudy, Matthews, and Houtz (15) found that 8 percent of the workers exposed to an average dust concentration of 5 million particles per cubic foot had asbestosis; 22 percent of the 17-million-particle group; and 57 percent of the 44-million-particle group had asbestosis.

THRESHOLD CONCENTRATION OF ASBESTOS DUST

From a practical standpoint, one of the most important results of a medical and engineering study such as this is the definition of safe working conditions in the industry under study. Ideally, a threshold concentration of dust should be the highest dust concentration that would not produce pneumoconiosis in originally healthy workmen during their entire working life. The chief difficulty in this study, as in most of the earlier studies of the Public Health Service, is that very few workmen are exposed for a long period of time to low concentrations of asbestos dust. Because of the importance of the problem, it seems to be desirable to use such data as are at hand to define tentative safe working conditions that may serve as standards for the guidance of factory managers and engineers until more complete data are available.

In an analysis of this kind the cases that occur at relatively low dust exposures are of especial significance because they indicate what the safe limits of exposure may be. These cases must be examined individually, as well as statistically, in order to be sure that there are no unusual features that would complicate statistical analysis. Below 2.5 million particles per cubic foot, the interpretation of table 43 offers no difficulties. None of the 39 persons exposed to dust concentrations below 2.5 million particles per cubic foot had a case of asbestosis, although, as a matter of fact, only 6 persons had been employed more than 5 years. Three doubtful cases of asbestosis fell in the range 2.5 to 4.9 million particles.

Because clean-cut cases of asbestosis were found only in dust concentrations exceeding 5 million particles per cubic foot, and because they were not found at lower dust concentrations, 5 million particles per cubic foot may be regarded tentatively as the threshold value for asbestos-dust exposure until better data are available.

In order to find out the levels to which asbestos-dust concentrations have been reduced in practice, an engineering study (7) was made in an asbestos textile factory in which dust control equipment has been recently installed. This shows that means are already available for reducing the dust exposure of a majority of asbestos textile workers to less than 5 million particles per cubic foot.

Published by AsbestosClaims.Law
asbestosclaims.law/library/18707/57